

Updated: 18 September 2019

[Print version](#)

# CEE 370 Environmental Engineering Principles

## Lecture #6 Environmental Chemistry IV: Thermodynamics, Equilibria, Acids-bases I

**Reading:** Mihelcic & Zimmerman, Chapter 3

Davis & Masten, Chapter 2  
Mihelcic, Chapt 3


David Reckhow

CEE 370 L#6

1

## Kinetics

- Base Hydrolysis of dichloromethane (DCM)
  - Forms chloromethanol (CM) and chloride



- Classic second order reaction (molecularity of 2)

$$\text{Rate} = k[\text{DCM}][\text{OH}^-] = -\frac{d[\text{DCM}]}{dt} = -\frac{d[\text{OH}^-]}{dt} = \frac{d[\text{CM}]}{dt} = \frac{d[\text{Cl}^-]}{dt}$$

- First order in each reactant, second order overall

David Reckhow

CEE 370 L#5


2

## Kinetic principles

- Law of Mass Action
  - For elementary reactions

$$aA + bB \xrightarrow{k} \text{products}$$

$$\text{rate} = kC_A^a C_B^b$$



where,

- $C_A$  = concentration of reactant species A, [moles/liter]
- $C_B$  = concentration of reactant species B, [moles/liter]
- $a$  = stoichiometric coefficient of species A
- $b$  = stoichiometric coefficient of species B
- $k$  = rate constant, [units are dependent on a and b]

David Reckhow
CEE 370 L#5
3

## Kinetics: zero order


- Reactions of order "n" in reactant "c"
 

$$\frac{dc}{dt} = -kc^n$$

---

- When  $n=0$ , we have a simple zero-order reaction
  - Example: biodegradation of 2,4-D
 

$$\frac{dc}{dt} = -k$$



Time (min)	Concentration
0	80
10	70
20	60
30	50
40	40
50	30
60	20
70	10

David Reckhow
CEE 370 L#5
4

## Kinetics: First Order

- When  $n=1$ , we have a simple first-order reaction
- This results in an "exponential decay"
  - Example: decay of  $^{137}\text{Cs}$  from Chernobyl accident

$$\frac{dc}{dt} = -kc^1$$

$$A \Rightarrow \text{products}$$

$c = c_0 e^{-kt}$

$k = 0.032 \text{ min}^{-1}$

David Reckhow
CEE 370 L#5
5

## Kinetics: First Order (cont.)

- This equation can be linearized
- good for assessment of "k" from data

$$\frac{dc}{dt} = -kc^1$$

$\ln c = \ln c_0 - kt$

Slope

$k = 0.032 \text{ min}^{-1}$

David Reckhow
CEE 370 L#5
6

## Kinetics: Second Order

$$\frac{dc}{dt} = -kc^2$$

- When  $n=2$ , we have a simple second-order reaction
 
$$A + A \Rightarrow \text{products}$$
- This is a reaction between two identical molecules
- More common to have different molecules reacting (see next slide)

Concentration

Time (min)

$k = 0.0015 \text{ L / mg / min}$

$c = c_o \frac{1}{1 + kc_o t}$

David Reckhow CEE 370 L#5 7


## Kinetics; Pseudo-1<sup>st</sup> Order

- When you have two different species reacting via 2<sup>nd</sup> order kinetics overall
 
$$A + B \Rightarrow \text{products}$$

$$\frac{dc_A}{dt} = \frac{dc_B}{dt} = -kc_A c_B$$
- If one reactant ( $c_B$ ) is present in great excess versus the other, the concentration of that one can be treated as constant and folded into the rate constant to get a **pseudo-first order** reaction
 
$$\frac{dc_A}{dt} = -k_{obs} c_A$$
 where  $k_{obs} = kc_B$

**From here you can use the first order equations**

David Reckhow CEE 370 L#5 8



## Half-lives

■ Time required for initial concentration to drop to half, i.e.,  $c = 0.5c_0$

■ For a zero order reaction:

$c = c_0 - kt$

$0.5c_0 = c_0 - kt_{1/2}$

$t_{1/2} = \frac{0.5c_0}{k}$

■ For a first order reaction:

$c = c_0 e^{-kt}$

$0.5c_0 = c_0 e^{-kt_{1/2}}$

$$t_{1/2} = \frac{\ln(2)}{k}$$

$$= \frac{0.693}{k}$$

Try example 3.5 & 3.6 in Mihelcic

David Reckhow CEE 370 L#5 20

## Temperature Effects

### Temperature Dependence

- Chemist's Approach: Arrhenius Equation
 

$$\frac{d(\ln k)}{dT_a} = \frac{E_a}{RT_a^2}$$

Activation energy

$$k_{T_a} = A e^{-E_a/RT_a}$$

Pre-exponential factor

$$k_{T_a} = k_{293^\circ K} e^{E_a(T_a - 293)/RT_a \cdot 293}$$

R = universal gas constant  
= 1.987 cal/°K/mole  
T<sub>a</sub> = absolute temp (°K)
- Engineer's Approach:
 

$$k_T = k_{20^\circ C} \theta^{T-20^\circ C}$$

Or more generally  
where T<sub>0</sub> is any  
"baseline" temperature

$$k_T = k_{T_0} \theta^{T-T_0}$$

Typical values:  
θ=1.02 to 1.15

David Reckhow 11


## Activity

- Activity is the "effective or apparent" concentration, which may be slightly different from the true "analytical" concentration
  - These two differ substantially in waters with high TDS, such as sea water.
- We identify these two as follows:
  - Curved brackets ({X}) indicate activity
  - Square brackets ([X]) indicate concentration
    - Usually this is molar concentration
    - This may also be used when we're not very concerned about the differences between activity and concentration

David Reckhow CEE 370 L#6 12

## Why the difference?

- Mostly long-range interactions between uninterested bystanders (chemical species that are not involved in the reaction) and the two dancers of interest (those species that are reacting)




David Reckhow CEE 370 L#6 13

## Activity & Ionic Strength

- Equilibrium quotients are really written for activities, not concentrations
 
$$K = \frac{\{C\}^c \{D\}^d}{\{A\}^a \{B\}^b}$$
- in most natural waters activities are nearly equal to the molar concentrations
 
$$\{A\} \approx [A]$$
- In saline waters, we must account for differences between the two
  - activity coefficients ( $f$  or  $\gamma$ ) are used for this
 
$$\{A\} = f_A [A]$$



$$\{A\} = \gamma_A [A]$$
  - Ionic Strength ( $I$  or  $\mu$ ) is used to determine the extent of correction
 
$$I \text{ or } \mu = \frac{1}{2} \sum C_i z_i^2$$

David Reckhow CEE 370 L#6 14



## Correlations for ionic strength

- $\mu$  vs. specific conductance: Russell Approximation
  - $\mu = 1.6 \times 10^{-5} \times K$  (in  $\mu\text{mho/cm}$ ) **Equ 3.3**
- $\mu$  vs. TDS: Langlier approximation
  - $\mu \sim 2.5 \times 10^{-5} \times \text{TDS}$  (in mg/L) **Equ 3.2**

David Reckhow
CEE 370 L#6
16



## Corrections to Ion Activity

Approximation	Equation	Applicable Range for I
Debye-Hückel	$\log f = -0.5z^2 \sqrt{I}$	$<10^{-2.3}$
Extended Debye-Hückel	$\log f = -0.5z^2 \frac{\sqrt{I}}{1+0.33a\sqrt{I}}$	$<10^{-1}$
Güntelberg	$\log f = -0.5z^2 \frac{\sqrt{I}}{1+\sqrt{I}}$	$<10^{-1}$ , solutions of multiple electrolytes
Davies	$\log f = -0.5z^2 \left( \frac{\sqrt{I}}{1+\sqrt{I}} - 0.2I \right)$	$<0.5$

From Stumm & Morgan, Table 3.3 (pg.103)

0.3, based on Mihelcic 17

David Reckhow CEE 370 L#6

## Kinetic model for equilibrium

- Consider a reaction as follows:
  - A + B = C + D
  - Since all reactions are reversible, we have two possibilities
 


$$A + B \xrightarrow{k_f} C + D$$

$$A + B \xleftarrow{k_b} C + D$$
  - The rates are:
 
$$r_f = k_f \{A\} \{B\} \quad r_b = k_b \{C\} \{D\}$$
  - And at equilibrium the two are equal,  $r_f = r_b$ 

$$k_f \{A\} \{B\} = k_b \{C\} \{D\}$$
  - We then define an equilibrium constant ( $K_{eq}$ )
 

$$K_{eq} \equiv \frac{k_f}{k_b} = \frac{\{C\} \{D\}}{\{A\} \{B\}}$$

David Reckhow CEE 370 L#6 18




## Kinetic model with moles

- In terms of molar concentrations, the rates are:
 
$$r_f = k_f [A] \gamma_A [B] \gamma_B \quad r_b = k_b [C] \gamma_C [D] \gamma_D$$
- And at equilibrium the two are equal,  $r_f = r_b$ 

$$k_f [A] \gamma_A [B] \gamma_B = k_b [C] \gamma_C [D] \gamma_D$$
- And solving for the equilibrium constant ( $K_{eq}$ )
 
$$K_{eq} \equiv \frac{k_f}{k_b} = \frac{[C] \gamma_C [D] \gamma_D}{[A] \gamma_A [B] \gamma_B} = \frac{[C][D] \left( \frac{\gamma_C \gamma_D}{\gamma_A \gamma_B} \right)}{[A][B] \left( \frac{\gamma_A \gamma_B}{\gamma_C \gamma_D} \right)}$$

David Reckhow CEE 370 L#6 19



## Equilibrium Chemistry

- Tells us what direction the reaction is headed in
- Doesn't tell us how fast the reaction is going (kinetics)
- Solving equilibrium problems
  - identify reactants and products
  - formulate equations
    - equilibrium equations
    - mass balance equations
    - electroneutrality equation
  - solve equations

David Reckhow CEE 370 L#6 20


## Temperature Effects on K

- Need  $\Delta H$  (enthalpy change)
  - $\Delta H < 0$ , exothermic (heat evolved)
  - $\Delta H > 0$ , endothermic (heat absorbed)
- The Van't Hoff Equation:
 

$$\log \frac{K_2}{K_1} = \frac{\Delta H^\circ (T_2 - T_1)}{2.303RT_2T_1}$$

  - recall that:
 

$$\Delta H^\circ = \sum \nu_i \Delta H_f^\circ$$



David Reckhow
CEE 370 L#6
21

## Acids & Bases


- pH of most mineral-bearing waters is 6 to 9. (fairly constant)
- pH and composition of natural waters is regulated by reactions of acids & bases
  - chemical reactions; mostly with minerals
    - carbonate rocks: react with  $\text{CO}_2$  (an acid)
      - $\text{CaCO}_3 + \text{CO}_2 = \text{Ca}^{+2} + 2\text{HCO}_3^-$
    - other bases are also formed:  $\text{NH}_3$ , silicates, borate, phosphate
    - acids from volcanic activity:  $\text{HCl}$ ,  $\text{SO}_2$
  - Biological reactions: photosynthesis & resp.
  - Sillen: Ocean is result of global acid/base titration

David Reckhow
CEE 370 L#6
22

## Acids & Bases (cont.)

- Equilibrium is rapidly established
  - proton transfer is very fast
- we call  $[H^+]$  the **Master Variable**
  - because Protons react with so many chemical species, affect equilibria and rates
- Strength of acids & bases
  - strong acids have a substantial tendency to donate a proton. This depends on the nature of the acid as well as the base accepting the proton (often water).

David Reckhow CEE 370 L#6 23



## Mathematical Expression of Acid/Base Strength

- Equilibrium constant
  - acids:  $HA = H^+ + A^-$ 
    - $HCl + H_2O = H_3O^+ + Cl^-$
    - $HCl = H^+ + Cl^-$
  - Bases:  $B + H_2O = BH^+ + OH^-$ 
    - $NH_3 + H_2O = NH_4^+ + OH^-$

$$K_a = \frac{[H^+][Cl^-]}{[HCl]} \approx 10^3$$

$$K_b = \frac{[NH_4^+][OH^-]}{[NH_3]} = 10^{-4.76}$$


David Reckhow CEE 370 L#6 25

## Thermodynamics & Equilibrium

- Classical Thermodynamics
- Ideal Gas Law
- Equilibrium Chemistry
  - Working with equations
  - Phase Transfer
  - Acids/Bases
    - The Carbonate System
  - Precipitation/Dissolution
  - Adsorption

} Review material on your own

David Reckhow CEE 370 L#7 26



## Enthalpy or heat of reaction

For a generic reaction:


$$aA + bB \rightarrow pP + rR$$

$$\Delta H^\circ_{rxn} = (p\Delta H^\circ_P + r\Delta H^\circ_R) + (a\Delta H^\circ_A + b\Delta H^\circ_B)$$

where,

A,B	=	reactant species
P,R	=	product species
a,b	=	stoichiometric coefficients of the reactants
p,r	=	stoichiometric coefficients of the products
$\Delta H^\circ_{A,B}$	=	enthalpy of reactants A and B
$\Delta H^\circ_{P,Q}$	=	enthalpy of products P and Q

David Reckhow CEE 370 L#7 27



## Enthalpy (cont.)


Or more generally:

$$\Delta H^\circ_{rxn} = \sum_{i=1}^n a_i \Delta H^\circ_{prod} - \sum_{j=1}^m a_j \Delta H^\circ_{rxtns}$$

where,

$a_i$	=	stoichiometric coefficient of product species
$i$		
$a_j$	=	stoichiometric coefficient of reactant
species j		
$\Delta H^\circ_i$	=	enthalpy of product species i, [kcal/mol]
$\Delta H^\circ_j$	=	enthalpy of reactant species j, [kcal/mol]

David Reckhow CEE 370 L#7 28



## Thermo- dynamic Constants

Table 4.2, pg. 57

Species	$\Delta G$ , kcal/mol	$\Delta H$ , kcal/mol
$\text{Ca}^{2+}(\text{aq})$	-132.18	-129.77
$\text{CaCO}_3(\text{s})$ , calcite	-269.78	-288.45
$\text{Ca}(\text{OH})_2(\text{s})$ , lime	-214.7	-235.7
$\text{CO}_2(\text{g})$	-94.26	-94.05
$\text{CO}_2(\text{aq})$	-92.31	-98.69
$\text{CO}(\text{aq})$	-126.2	-161.6
$\text{Cl}(\text{aq})$	-31.3	-40.0
$\text{Cl}_2(\text{aq})$	-19.1	-28.9
$\text{CO}(\text{aq})$	-8.8	-25.6
$\text{H}(\text{aq})$	0	0
$\text{HNO}_3(\text{aq})$	-26.6	-25.0
$\text{NO}(\text{aq})$	-26.6	-49.5
$\text{HCO}(\text{aq})$	-19.1	-28.9
$\text{CH}(\text{aq})$	-37.6	-55.0
$\text{O}(\text{aq})$	3.9	-267
$\text{H}_2\text{O}(\text{l})$	-68.7	-68.3
$\text{H}(\text{g})$	0	0
$\text{O}(\text{g})$	0	0

David Reckhow
CEE 370 L#7
29

### Thermodynamic Constants for Species of Importance in Water Chemistry (Table 3-1 from Snoeyink & Jenkins) Part I

Species	$\Delta \overline{H}_f^\circ$ kcal/mole	$\Delta \overline{G}_f^\circ$ kcal/mole	Species	$\Delta \overline{H}_f^\circ$ kcal/mole	$\Delta \overline{G}_f^\circ$ kcal/mole
$\text{Ca}^{+2}(\text{aq})$	-129.77	-132.18	$\text{CO}_3^{-2}(\text{aq})$	-161.63	-126.22
$\text{CaCO}_3(\text{s})$ , calcite	-288.45	-269.78	$\text{CH}_3\text{COO}^-$ , acetate	-116.84	-89.0
$\text{CaO}(\text{s})$	-151.9	-144.4	$\text{H}^+(\text{aq})$	0	0
$\text{C}(\text{s})$ , graphite	0	0	$\text{H}_2(\text{g})$	0	0
$\text{CO}_2(\text{g})$	-94.05	-94.26	$\text{Fe}^{+2}(\text{aq})$	-21.0	-20.30
$\text{CO}_2(\text{aq})$	-98.69	-92.31	$\text{Fe}^{+3}(\text{aq})$	-11.4	-2.52
$\text{CH}_4(\text{g})$	-17.889	-12.140	$\text{Fe}(\text{OH})_3(\text{s})$	-197.0	-166.0
$\text{H}_2\text{CO}_3(\text{aq})$	-167.0	-149.00	$\text{Mn}^{+2}(\text{aq})$	-53.3	-54.4
$\text{HCO}_3^-(\text{aq})$	-165.18	-140.31	$\text{MnO}_2(\text{s})$	-124.2	-111.1

Conversion: 1kcal = 4.184 kJ

David Reckhow
CEE 370 L#7
30

## Thermodynamic Constants for Species of Importance in Water Chemistry (Table 3-1 from Snoeyink & Jenkins) Part II

Species	$\Delta \overline{H}_f^\circ$ kcal/mole	$\Delta \overline{G}_f^\circ$ kcal/mole	Species	$\Delta \overline{H}_f^\circ$ kcal/mole	$\Delta \overline{G}_f^\circ$ kcal/mole
Mg <sup>2+</sup> (aq)	-110.41	-108.99	O <sub>2</sub> (g)	0	0
Mg(OH) <sub>2</sub> (s)	-221.00	-199.27	OH <sup>-</sup> (aq)	-54.957	-37.595
NO <sub>3</sub> <sup>-</sup> (aq)	-49.372	-26.43	H <sub>2</sub> O (g)	-57.7979	-54.6357
NH <sub>3</sub> (g)	-11.04	-3.976	H <sub>2</sub> O (l)	-68.3174	-56.690
NH <sub>3</sub> (aq)	-19.32	-6.37	SO <sub>4</sub> <sup>2-</sup>	-216.90	-177.34
NH <sub>4</sub> <sup>+</sup> (aq)	-31.74	-19.00	HS (aq)	-4.22	3.01
HNO <sub>3</sub> (aq)	-49.372	-26.41	H <sub>2</sub> S(g)	-4.815	-7.892
O <sub>2</sub> (aq)	-3.9	3.93	H <sub>2</sub> S(aq)	-9.4	-6.54

Conversion: 1kcal = 4.184 kJ

David Reckhow

CEE 370 L#7

31

## Gibbs Free Energy

$$\Delta G = \Delta H - T\Delta S$$

where,


$\Delta G$ [kcal/mol]	=	Gibbs free energy,
$\Delta H$	=	enthalpy, [kcal/mol]
$\Delta S$	=	entropy, [kcal/K-mol]
T	=	temperature, [K]

David Reckhow

CEE 370 L#7

32






## Gibbs Free Energy

- Combines enthalpy and entropy
  - 1st and 2nd laws of thermodynamics
- Determines whether a reaction is favorable or spontaneous
- Practical form is based on an arbitrary datum
  - the pure and most stable form of each element at standard state

$$\Delta G^{\circ} = \Delta H^{\circ} - T\Delta S^{\circ}$$

David Reckhow CEE 370 L#7 33



## Reactions under Standard State Conditions


$$aA + bB \rightarrow pP + rR$$

$$\Delta G^{\circ}_{rxn} = (p\Delta G^{\circ}_P + r\Delta G^{\circ}_R) - (a\Delta G^{\circ}_A + b\Delta G^{\circ}_B)$$

where,

- A,B = reactant species
- P,R = product species
- a,b = stoichiometric coefficients of the reactants
- p,r = stoichiometric coefficients of the reactants
- $\Delta G^{\circ}_{A,B}$  = Gibbs free energy of reactants A and B
- $\Delta G^{\circ}_{P,Q}$  = Gibbs free energy of reactants P and Q

David Reckhow CEE 370 L#7 34




## General Equation

$$\Delta G^{\circ}_{rxn} = \left( \sum_{i=1}^n a_i \Delta G^{\circ}_i \right)_{prod} - \left( \sum_{j=1}^m a_j \Delta G^{\circ}_j \right)_{rxnts}$$

where,

- $a_i$  = stoichiometric coefficient of product species i
- $a_j$  = stoichiometric coefficient of reactant species j
- $\Delta G^{\circ}_i$  = standard Gibbs free energy of product species i, [kcal/mol]
- $\Delta G^{\circ}_j$  = standard Gibbs free energy of reactant species j, [kcal/mol]

David Reckhow CEE 370 L#7 35



## Non-standard state conditions

$$\Delta G = \Delta G^{\circ} + RT \ln \frac{P^p R^r}{A^a B^b}$$

And we define:

$$Q = \frac{P^p R^r}{A^a B^b}$$

At equilibrium,  $\Delta G=0$ , and  $Q=K_{eq}$

$$\Delta G^{\circ} = -RT \ln K_{eq}$$

**End** equilibria

David Reckhow CEE 370 L#7 36

## What are standard state conditions?

Parameter	Standard State or Condition
Temperature	25°C
Gas	1 atm
Solid	Pure solid
Liquid	Pure liquid
Solution	1 M
Element*	0


David Reckhow
CEE 370 L#7
37

## Roller Coaster Analogy

Fig. 3-8 in Mihelcic

A reaction with a negative  $\Delta G$  will proceed from reactants to products. A reaction with a positive  $\Delta G$  will proceed from products to reactants (backwards). A reaction in which  $\Delta G$  is zero is at equilibrium


David Reckhow
CEE 370 L#7
38



## Temperature effects

- Increases in temperature cause:
  - A. Increases in solubility of salts
  - B. Decreases in solubility of salts
  - C. Depends on the salt


David Reckhow CEE 370 L#6 39



## Temperature effects

- Increases in temperature cause:
  - A. Increases in solubility of dissolved gases
  - B. Decreases in solubility of dissolved gases
  - C. Depends on the gas

David Reckhow CEE 370 L#6 40




## Lime Example

Lime, or calcium hydroxide, is commonly used to improve sedimentation processes or to precipitate toxic metals. A 500 L reaction vessel contains an excess amount of solid calcium hydroxide. The solution contains  $10^{-8}$  M of  $\text{Ca}^{2+}$  and  $10^{-8}$  M of  $\text{OH}^-$ . Is the lime still dissolving, is the solution at equilibrium, or is lime precipitating?

$$\text{Ca(OH)}_2(\text{s}) \rightarrow \text{Ca}^{2+} + 2\text{OH}^-$$

Example 4.6 from Ray

David Reckhow CEE 370 L#7 41



## Solution to Lime example


First calculate the Gibbs Free Energy under standard state conditions

$$\Delta G^\circ_{\text{rxn}} = (\Delta G^\circ_{\text{Ca}^{2+}} + 2\Delta G^\circ_{\text{OH}^-}) - \Delta G^\circ_{\text{Ca(OH)}_2}$$

$$\Delta G^\circ_{\text{rxn}} = [(-132.3 \text{ kcal/mol}) + 2 \times (-37.6 \text{ kcal/mol}) - (-214.7 \text{ kcal/mol})]$$

$$\Delta G^\circ_{\text{rxn}} = +7.22 \text{ kcal/mol}$$

David Reckhow CEE 370 L#7 42



## Solution (cont.)

Now we need to adapt to actual conditions:

$$\Delta G = \Delta G^\circ + RT \ln \frac{P^p R^r}{A^a B^b} \quad \text{Ca(OH)}_2(\text{s}) \rightarrow \text{Ca}^{2+} + 2 \text{OH}^-$$


$$\Delta G_{\text{rxn}} = \Delta G^\circ + RT \ln [\text{Ca}^{2+}][\text{OH}^-]^2$$

$$\Delta G_{\text{rxn}} = \left(7.22 \frac{\text{kcal}}{\text{K-mol}}\right) + \left(1.987 \frac{\text{cal}}{\text{K-mol}}\right) \times \left(\frac{\text{kcal}}{1000 \text{ cal}}\right) \times (298 \text{ K}) \times \left(\ln (10^{-8})(10^{-8})^2\right)$$

$$\Delta G_{\text{rxn}} = -13.2 \text{ kcal/mol}$$

Since  $\Delta G$  is negative, the reaction will proceed as written


David Reckhow CEE 370 L#7 43



## Kinetics & Equilibrium

- Kinetics
  - Rates of reactions
  - Dynamic, complex
  - Best approach for slow reactions
    - Oxidation reactions, biochemical transformations, photochemical reactions, radioactive decay
- Equilibrium: thermodynamics
  - Final stopping place
  - Static, simple
  - Best approach for fast reactions
    - Acid/base reaction, complexation, some phase transfer

David Reckhow CEE 370 L#7 44




## DO Example

A polluted stream with a temperature of 25°C has a dissolved oxygen concentration of 4 mg/L. Use Gibbs free energy to determine if oxygen from the atmosphere is dissolving into the water, the oxygen is at equilibrium, or oxygen from the stream is going into the atmosphere.

$$\text{O}_2(\text{aq}) \rightarrow \text{O}_2(\text{g})$$

Example 4.7 from Ray

David Reckhow CEE 370 L#7 45



## Solution to DO example


$$\Delta G^\circ_{\text{rxn}} = \Delta G^\circ_{\text{O}_2(\text{g})} - \Delta G^\circ_{\text{O}_2(\text{aq})} = \left(0 \frac{\text{kcal}}{\text{mol}}\right) - \left(-3.9 \frac{\text{kcal}}{\text{mol}}\right)$$

$$\Delta G^\circ_{\text{rxn}} = 3.9 \frac{\text{kcal}}{\text{mol}}$$

$$\Delta G = \Delta G^\circ + RT \ln \frac{P_{\text{O}_2}}{[\text{O}_2(\text{aq})]}$$

$$[\text{O}_2(\text{aq})] = 4 \frac{\text{mg O}_2}{\text{L}} \times \frac{\text{g O}_2}{1000 \text{ mg O}_2} \times \frac{\text{mol O}_2}{32 \text{ g O}_2} = 1.25 \times 10^{-4} \text{ M}$$

David Reckhow CEE 370 L#7 46




## Solution (cont.)

$$\Delta G = -3.9 \frac{\text{kcal}}{\text{mol}} \times \frac{1000 \text{ cal}}{\text{kcal}} + \left(1.987 \frac{\text{cal}}{\text{K-mol}}\right) (298 \text{ K}) \ln\left(\frac{0.209 \text{ atm}}{1.25 \times 10^{-4} \text{ M}}\right)$$
$$\Delta G = 491 \frac{\text{cal}}{\text{mol}} > 0$$

Since  $\Delta G$  is positive, the reaction will proceed in the reverse direction as written. From the atmosphere to the water.

David Reckhow CEE 370 L#7 47



■ To next lecture

David Reckhow CEE 370 L#6 48