

Updated: 6 November 2019 [Print version](#)

# CEE 370 Environmental Engineering Principles

Lecture #25  
Water Quality Management III:  
Lakes & toxic models

[Reading: Mihelcic & Zimmerman, Chapter 7 & 3.10](#)  
[Reading: Davis & Cornwall, Chapt 5-4](#)  
[Reading: Davis & Masten, Chapter 5-6 & 9-4](#)

David Reckhow CEE 370 L#25 1

## Lake Pollution

- Percent impaired by pollutant
- Percent impaired by sources

From Masters, section 5.4

**Pollutants**

Pollutant	Total
Metals	47
Nutrients	40
Organic enrichment/DO	24
Siltation	22
Priority organic chemicals	20

(a) By pollutant

**Pollutant sources**

Source	Total
Agriculture	56
Urban runoff/storm sewers	24
Hydrologic/habitat modification	23
Municipal point sources	21
Onsite wastewater disposal	16

David Reckhow

## Lake life cycle

- Succession in lakes
  - Oligotrophic
  - Mesotrophic
  - Eutrophic
  - Other
    - Dystrophic
    - Hyper-eutrophic

Fertiliser run-off

1. Algae grow fast, using up lots of oxygen and blocking sunlight
2. Aquatic plants begin to die
3. Dead matter provides food for microbes ...
4. ... increasing the competition for oxygen
5. Water becomes deoxygenated - fish die

David Reckhow


## Lake Eutrophication

- As lakes age, they become more productive
  - Natural processes: natural Eutrophication
  - Pollutant loading: cultural Eutrophication
- Limiting nutrient
  - Liebig's law of minimum
  - Redfield Ratio
    - C:N:P in most phytoplankton is 106:16:1
    - When  $P < 16 \cdot N$ , it limits growth

David Reckhow

CEE 370 L.#25

4



## Estrogen lake study

### Collapse of a fish population after exposure to a synthetic estrogen

Karen A. Kidd<sup>1\*</sup>, Paul J. Blanchfield<sup>2\*</sup>, Kenneth H. Mills<sup>2\*</sup>, Vince P. Palace<sup>2\*</sup>, Robert E. Evans<sup>2\*</sup>, James M. Lazorchak<sup>2</sup>, and Robert W. Flick<sup>2</sup>

<sup>1</sup>Fisheries and Oceans Canada, Freshwater Institute, 501 University Crescent, Winnipeg, Manitoba, Canada R3T 2N6; and <sup>2</sup>Molecular Indicators Research Branch, United States Environmental Protection Agency, 26 West Martin Luther King Drive, Cincinnati, OH 45268

Edited by Deborah Swackhamer, University of Minnesota, Minneapolis, MN, and accepted by the Editorial Board March 29, 2007 (received for review October 27, 2006)

Municipal wastewaters are a complex mixture containing estrogens and estrogen mimics that are known to affect the reproductive health of wild fishes. Male fishes downstream of some wastewater outfalls produce vitellogenin (VTG) (a protein normally synthesized by females during oocyte maturation) and early-stage eggs in their testes, and this feminization has been attributed to the presence of estrogenic substances such as natural estrogens [estrone or 17 $\beta$ -estradiol (E2)], the synthetic estrogen used in birth-control pills [17 $\alpha$ -ethynylestradiol (EE2)], or weaker estrogen mimics such as nonylphenol in the water. Despite widespread evidence that male fishes are being feminized, it is not known whether these low-level, chronic exposures adversely impact the sustainability of wild populations. We conducted a 7-year, whole-lake experiment at the Experimental Lakes Area (ELA) in northwestern Ontario, Canada, and showed that chronic exposure of fathead minnow (*Pimephales promelas*) to low concentrations (5–6 ng L<sup>-1</sup>) of the potent 17 $\alpha$ -ethynylestradiol led to feminization of males through the production of vitellogenin mRNA and protein, impacts on gonadal development as evidenced by intersex in males and altered oogenesis in females, and, ultimately, a near extinction of this species from the lake. Our observations demonstrate that the concentrations of estrogens and their mimics observed in freshwaters can impact the sustainability of wild fish populations.

present (9) and has been linked to the feminization of male fishes in rivers receiving municipal wastewater (4, 6). Despite growing documentation on the feminization of male fishes in waterways receiving municipal effluents, a critical question in the field of endocrine disruption research is whether these low-level, chronic exposures adversely impact wild populations (13). Although laboratory studies have shown decreased reproductive success of fish exposed to <1–5 ng L<sup>-1</sup> of EE2 (14, 15), it is unknown whether this response would be observed in wild populations and whether it would result in a subsequent decline in abundances. To assess the ecological risk posed by this class of compounds, we must understand population-level effects of estrogens and their mimics on aquatic organisms.

The fathead minnow (*Pimephales promelas*) is a common species in North America, and its range extends from the southern United States to northern Canada (16). It is an important food source for numerous game fish species, such as lake trout (*Salvelinus namaycush*), walleye (*Sander vitreus*), and northern pike (*Esox lucius*). In the lakes used for this study, fathead minnow have a lifespan of  $\approx$ 4 years, but few individuals live past 2 years of age (17). Asynchronous spawning starts in early summer and extends for a period of  $\approx$ 2 months; multiple females will typically spawn in the nest of a single male, who will

## Concern over drinking water

- Drugs?



David Reckhow

CEE 370 L#25

7


## Organic Compounds: Types?

- Natural Compounds
  - Fulvics
  - Proteins, carbohydrates, etc
- Domestic WW Organics
- Industrial Synthetic Organics
  - Plasticizers: phthalates
  - solvents: tetrachloroethylene
  - waxes: chlorinated parafins
  - others: PCB's
- Hydrocarbons & oil derivatives
  - includes products of combustion: PAH's
- Agricultural Chemicals
  - pesticides: DDT, kepone, mirex
- Pharmaceuticals, etc
  - Anti-epileptics
  - Beta-blockers
  - X-ray contrast media
  - antibiotics
- Home & Personal Care Products
  - triclosan
  - Musks, flame retardants
- Endocrine Disrupters
  - Steroidal estrogens
- Natural process byproducts
  - Conjugated pharmaceuticals
- Engineered process byproducts
  - disinfection byproducts, etc

David Reckhow

CEE 370 L#25


8



## Pollutant loss in lakes

- Photolysis
  - Destruction by solar light energy
- Biodegradation
  - Metabolism by microorganism
- Hydrolysis
  - Chemical decomposition
- Volatilization
  - Loss to the atmosphere
- Adsorption and settling
  - Loss to particles that end up buried in sediments



David Reckhow CEE 370 L.#25 9




## Photolysis

- Chemical breakdown initiated by light energy
- two types
  - direct photolysis
  - sensitized (or indirect) photolysis
- Several steps
  - some solar light reaches water surface
  - some of this light penetrates to the solute
  - some of this is absorbed by the solute
  - some of absorbed light is capable of causing a reaction

David Reckhow CEE 370 L.#25 10






## Biotransformation

- Microbially mediated transformation of organic and inorganic contaminants
- Biochemical processes:
  - Metabolism: toxicant is used for synthesis or energy
  - Cometabolism: not "used", but transformed anyway
- Chemical Effects:
  - Detoxication: Toxic to Non-toxic
    - mineralization
  - Activation: Non-toxic to Toxic

David Reckhow CEE 370 L#25 13

Refer to lecture #17




## Bio kinetics

- Michaelis-Menten equation:
  - $\mu_{\max}$  = maximum growth rate ( $\text{yr}^{-1}$ )
  - $X$  = microbial biomass ( $\# \text{cells}/\text{m}^3$ )
  - $Y$  = yield coefficient (cells produced per mass toxicant removed,  $\# \text{cells}/\mu\text{g}$ )
  - $k_s$  = half-saturation constant ( $\mu\text{g}/\text{m}^3$ )
  - $k_b$  = rate of biotransformation ( $\text{yr}^{-1}$ )
$$k_m = \frac{\mu_{\max} X}{Y(k_s + c)}$$
- If  $c \ll k_s$ , then:
 
$$k_m = \frac{\mu_{\max} X}{Y k_s} = k_{m2} X$$

David Reckhow CEE 370 L#25 14

## Bio kinetics (cont.)

- Wide environmental range
  - phenol:  $k_m = 4.0 \text{ d}^{-1}$
  - diazinon:  $k_m = 0.016 \text{ d}^{-1}$
- Temperature correction
  - $\theta = 1.04 - 1.095$



$$(k_m)_T = (k_m)_{20} \theta^{T-20}$$


David Reckhow CEE 370 L#25 15

## Hydrolysis

- Reaction with water and its constituents
  - $\text{H}_2\text{O}$   $k_h = k_n$
  - $\text{OH}^-$   $k_h = k_b [\text{OH}^-]$
  - $\text{H}^+$   $k_h = k_a [\text{H}^+]$
- Autodissociation  $K_w = [\text{OH}^-] [\text{H}^+]$
- Combining:  $k_h = k_b [\text{OH}^-] + k_n + k_a [\text{H}^+]$ 
  - or:  $k_h = k_b \frac{K_w}{10^{-\text{pH}}} + k_n + k_a 10^{-\text{pH}}$

David Reckhow CEE 370 L#25 16





## Two film model

- Flux from the bulk liquid to the interface
 
$$J_l = K_l(c_i - c)$$
- Flux from the interface of the bulk gas
 
$$J_g = \frac{K_g}{RT_a}(p_g - p_i)$$

Mass transfer velocities (m/d)

- And the K's are related to the molecular diffusion coefficients by:
 
$$K_l = \frac{D_l}{z_l} \qquad K_g = \frac{D_g}{z_g}$$

18

## Whitman's 2 film model (cont.)

- According to Henry's law:  $p_i = H_e c_i$
- And relating this back to the bulk concentration
 
$$p_i = H_e \left( \frac{J_l}{K_l} + c \right)$$
- now solving and equating the fluxes, we get:
 
$$\frac{1}{v_v} = \frac{1}{K_l} + \frac{RT_a}{H_e K_g}$$

The net transfer velocity across the air-water interface (m/d)

David Reckhow CEE 370 L#25 19

## Whitman's 2 film model (cont.)

- Which can be rewritten as:
 
$$v_v = K_l \frac{H_e}{H_e + RT_a \left( \frac{K_l}{K_g} \right)}$$
- Now, applying it to toxicants
  - $p_g \approx 0$
- And converting to the appropriate units:
 
$$k_v = \frac{v_v}{H}$$

David Reckhow CEE 370 L#25 20

## Volatilization: Parameter estimation

- Liquid film mass transfer coefficient ( $d^{-1}$ )
 
$$K_l = K_{l,O_2} \left( \frac{32}{M} \right)^{0.25}$$


Compound molecular weight
- Gas film mass transfer coefficient ( $d^{-1}$ )
 
$$K_g = 168U_w \left( \frac{18}{M} \right)^{0.25}$$

Wind velocity (mps)

David Reckhow CEE 370 L#25 21

## Toxics Model: CSTR with sediments

- Internal Transport Processes (between compartments)
  - dissolved: diffusion
  - particulate: settling, resuspension & burial
- Expressed as velocities (e.g., m/yr)



Water (1)

Mixed Sediments (2)

Deep Sediments (3)

Diffusion ( $v_d$ )

Settling ( $v_s$ )

Resuspension ( $v_r$ )

Burial ( $v_b$ )

David Reckhow CEE 370 L#25

## Sorption

Functionally identical to M&Z equ# 3.32

$K = \frac{q}{C}$

■ Linear Isotherm  $K = \frac{C_{particulate}}{C_{dissolved}}$

See M&Z, section 3.10

$c'_d \longleftrightarrow c_p$

$\text{Hexagon} \xrightleftharpoons[M_s k_{ad}]{k_{de}} \text{Hexagon} \text{ Particle}$

$k_s = v_s \left( \frac{K}{1+K} \right)$

Particle can then settle

$v_s$

23


David Reckhow CEE 370 L#25

Also in Lecture #16 biomagnification & Lecture #22 on groundwater retardation

## Octanol:water partitioning

- 2 liquid phases in a separatory funnel that don't mix
  - octanol
  - water
- Add contaminant to flask
- Shake and allow contaminant to reach equilibrium between the two
- Measure concentration in each ( $K_{ow}$  is the ratio)
- Correlate to environmental K

$K = fn(K_{ow})$




24

David Reckhow CEE 370 L#25


From Lecture #16

## Bioaccumulation


- Mercury in food chain
  - Data from Onondaga Lake



Biomass  
(box size)



Concentration  
(Shading)




Walleye (piscivore)	2.27 µg/g
Smallmouth Bass (piscivore)	0.91 µg/g
Gizzard Shad (planktivore)	0.23 µg/g
Zooplankton	$9.1 \times 10^{-4}$ µg/g
Phytoplankton	$5.6 \times 10^{-6}$ µg/g

David Reckhow

From Lecture #16

## Lab to Field

- Octanol water partition coefficients and bioconcentration factors



David Reckhow

CEE 370 L#25

26

## Completely-mixed lake or CMFR

- Often useful to assume perfect mixing
  - same concentration throughout system

*Accumulation = loading – outflow – reaction – settling*

Mass Balance

David Reckhow
CEE 370 L#25
27

## Other Terms in the Mass Balance

- Outflow *Outflow = Qc*
- Reaction *Reaction = kM = kVc*
- Settling *Settling = vA<sub>s</sub>c*  
*= k<sub>s</sub>Vc*

Note HW#6,  
problem 2

Since:

$k_s = v/H$

$V = A_s H$

David Reckhow
CEE 370 L#25
28

## Combining all terms:

$$V \frac{dc}{dt} = W(t) - Qc - kVc - vA_s c$$

- Units for each term: mass/time
- Dependent variable:  $c$
- Independent variable:  $t$
- Forcing function:  $W(t)$ , the way in which the external world "forces" the system
- Parameters:  $V, Q, k, v, A_s$

David Reckhow CEE 370 L#25 29

## Steady State Case

**With Settling**

- mass balance  $V \frac{dc}{dt} = 0 = W(t) - Qc - kVc - vA_s c$
- solution  $c = \frac{W}{Q + kV + vA_s}$  or  $c = \frac{W}{a}$
- assimilation factor  
 Where:  $a = Q + kV + vA_s$   
 The assimilation or "cleansing" factor

David Reckhow CEE 370 L#25 30

### Simple lake model without settling

[accumulation] = [loadings] ± [transport] ± [reactions]

$$V \frac{dc}{dt} = W(t) - Qc - kVc^n$$

For a 1st order reaction (n=1):

$$\frac{dc}{dt} + \alpha c = \frac{W(t)}{V}$$

Where:  $\alpha = \frac{Q}{V} + k$

Steady State Solution:  $c = \frac{\bar{W}}{Q + kV}$

David Reckhow CEE 370 L#25 31

### Lake Model: Steady State Example

A lake has the following characteristics:


*Volume* = 50,000 m<sup>3</sup>  
*Mean Depth* = 2 m  
*Inflow* = *Outflow* = 7500 m<sup>3</sup>d<sup>-1</sup>  
*Temperature* = 25°C

The lake receives the input of a pollutant from three sources: a factory discharge of 50 kg d<sup>-1</sup>, a flux from the atmosphere of 0.6 g m<sup>-2</sup> d<sup>-1</sup>, and the inflow stream that has a concentration of 10 mg/L. If the pollutant decays at the rate of 0.25/d at 20°C

- a. compute the assimilation factor
- b. steady state concentration
- c. show breakdown for each term

David Reckhow CEE 370 L#25 32





## Lake Model: Solution


First correct the decay rate for temperature

$$k = 0.25\theta^{25-20} = 0.25(1.05)^{25-20} = 0.319d^{-1}$$

Now the assimilation factor

$$a = Q + kV = 7500 + 0.319(50,000) = 23,454m^3d^{-1}$$

David Reckhow CEE 370 L#25 33



## Lake Model: Solution (cont.)

The surface area of the lake is:

$$A_s = \frac{V}{H} = \frac{50,000}{2} = 25,000m^2$$

The atmospheric and inflow load is then:

$$W_{atmosphere} = JA_s = 0.6(25,000) = 15,000g/d$$

$$W_{inflow} = 7500(10) = 75,000g/d$$

Combining all loads:

$$W = W_{factory} + W_{atmosphere} + W_{inflow} = 50,000 + 15,000 + 75,000 = 140,000g/d$$

David Reckhow CEE 370 L#25 34

## Lake Model: Solution (cont.)

And finally, the concentration:

$$c = \frac{W}{a}$$
$$= \frac{140,000 \text{ g} / \text{d}}{23,454 \text{ m}^3 / \text{d}}$$
$$= 5.97 \text{ mg} / \text{L}$$

David Reckhow CEE 370 L#25 35



## Dimictic Lakes


### Lake Stratification

Epilimnion

Thermocline

David Reckhow CEE 370 L#25 36





■ To next lecture

David Reckhow CEE 370 L#25 40