



Updated: 24 April 2020 [Print version](#)

CEE 680: Water Chemistry

Lecture #46
Redox Chemistry: Basic Calculations
(Stumm & Morgan, Chapt.8)
Benjamin; Chapter 9

David Reckhow CEE 680 #47 1





AGU PUBLICATIONS

Global Biogeochemical Cycles

RESEARCH ARTICLE
10.1002/2017GB005749

Key Points:

- Fe has increased significantly in 28% of the 340 studied waters by on average 64% between 1990 and 2013
- Most increasing trends and the highest rate of change were found in the boreal region of northern Europe
- Hydrological changes may have contributed to the observed Fe trends. Trends may also be linked to declining atmospheric S deposition

Correspondence to:
C. Björnerås,
caroline.bjorneras@bioluse.se

Citation:
Björnerås, C., Weyhenmeyer, G. A., Evans, C. D., Gessner, M. O., Grossart, H.-P., Kangur, K., ... Kritzberg, E. S. (2017). Widespread increases in iron concentration in European and North American freshwaters. *Global Biogeochemical Cycles*, 31, 1488–1500.

Widespread Increases in Iron Concentration in European and North American Freshwaters

C. Björnerås¹ , G. A. Weyhenmeyer² , C. D. Evans³ , M. O. Gessner⁴ , H.-P. Grossart⁵ , K. Kangur⁶ , I. Kokorite⁷, P. Kortelainen⁸ , H. Laudon⁹ , J. Lehtoranta⁸, N. Lottig¹⁰ , D. T. Monteith¹¹, P. Nöges¹², T. Nöges¹², F. Oulehle¹³ , G. Riise¹⁴ , J. A. Rusak¹⁵ , A. Räike⁸, J. Sire¹⁶, S. Sterling¹⁷ , and E. S. Kritzberg¹

¹Aquatic Ecology, Department of Biology, Lund University, Lund, Sweden, ²Ecology and Genetics/Limnology, Evolutionary Biology Centre, Uppsala University, Uppsala, Sweden, ³Centre for Ecology and Hydrology, Environment Centre Wales, Bangor, UK, ⁴Department of Experimental Limnology, Leibniz Institute of Freshwater Ecology and Inland Fisheries (IGB), Stechlin, Germany and Department of Ecology, Berlin Institute of Technology (TU Berlin), Berlin, Germany, ⁵Department Experimental Limnology, Leibniz Institute of Freshwater Ecology and Inland Fisheries, Stechlin, Germany and Institute of Biochemistry and Biology, Potsdam University, Potsdam, Germany, ⁶Centre for Limnology, Institute of Agricultural and Environmental Sciences, Estonian University of Life Sciences, Rannu, Estonia, ⁷Institute of Biology, University of Latvia, Riga, Latvia, ⁸Finnish Environment Institute, Helsinki, Finland, ⁹Department of Forest Ecology and Management, Swedish University of Agricultural Sciences, Umeå, Sweden, ¹⁰Trout Lake Station, Center for Limnology, University of Wisconsin-Madison, Wisconsin, USA, ¹¹NERC Centre for Ecology and Hydrology, Lancaster Environment Centre, Lancaster, UK, ¹²Centre for Limnology, Institute of Agricultural and Environmental Sciences, Estonian University of Life Sciences, Tartu, Estonia, ¹³Czech Geological Survey, Klárov 3, Prague, Czech Republic and Global Change Research Institute CAS, Brno, Czech Republic, ¹⁴Norwegian University of Life Sciences (NMBU), Ås, Norway, ¹⁵Dorset Environmental Science Centre, Ontario Ministry of the Environment and Climate Change, Dorset, Ontario, Canada, ¹⁶Inland Waters Division, Latvian Environment Geology and Meteorology Centre, Riga, Latvia, ¹⁷Hydrology Research Group, Department of Earth Sciences, Dalhousie University, Halifax, Nova Scotia, Canada

Question for the class

- Which is the strongest chemical oxidant?
 1. Chlorine
 2. Chloramines
 3. Permanganate
 4. Ozone
 5. Chlorine Dioxide
 6. Hydrogen Peroxide

David Reckhow

CEE 680 #46

5

Standard Half Cell Potentials
for Some Oxidation Reactions that Can Occur During Drinking Water Treatment

Oxidant	Reduction half-reaction	E°_{red} , volts
Ozone	$\frac{1}{2}\text{O}_3(\text{aq}) + \text{H}^+ + \text{e}^- \rightarrow \frac{1}{2}\text{O}_2(\text{aq}) + \text{H}_2\text{O}$	2.04
Hydrogen Peroxide	$\frac{1}{2}\text{H}_2\text{O}_2 + \text{H}^+ + \text{e}^- \rightarrow \text{H}_2\text{O}$	1.78
Permanganate	$\frac{1}{3}\text{MnO}_4^- + \frac{4}{3}\text{H}^+ + \text{e}^- \rightarrow \frac{1}{3}\text{MnO}_{2(\text{s})} + \frac{2}{3}\text{H}_2\text{O}$	1.68
Chlorine Dioxide	$\text{ClO}_2 + \text{e}^- \rightarrow \text{ClO}_2^-$	1.15
Hypochlorous Acid	$\frac{1}{2}\text{HOCl} + \frac{1}{2}\text{H}^+ + \text{e}^- \rightarrow \frac{1}{2}\text{Cl}^- + \frac{1}{2}\text{H}_2\text{O}$	1.49
Hypochlorite Ion	$\frac{1}{2}\text{OCl}^- + \text{H}^+ + \text{e}^- \rightarrow \frac{1}{2}\text{Cl}^-$	0.90
Hypobromous acid	$\frac{1}{2}\text{HOBr} + \frac{1}{2}\text{H}^+ + \text{e}^- \rightarrow \frac{1}{2}\text{Br}^- + \frac{1}{2}\text{H}_2\text{O}$	1.33
Monochloramine	$\frac{1}{2}\text{NH}_2\text{Cl} + \text{H}^+ + \text{e}^- \rightarrow \frac{1}{2}\text{Cl}^- + \frac{1}{2}\text{NH}_4^+$	1.40
Dichloramine	$\frac{1}{4}\text{NHCl}_2 + \frac{3}{4}\text{H}^+ + \text{e}^- \rightarrow \frac{1}{2}\text{Cl}^- + \frac{1}{4}\text{NH}_4^+$	1.34
Oxygen	$\frac{1}{2}\text{O}_2(\text{aq}) + \text{H}^+ + \text{e}^- \rightarrow \frac{1}{2}\text{H}_2\text{O}$	1.27

David Reckhow

CEE 680 #46

6

Standard Half Cell Potentials for Some Oxidation Reactions that Can Occur During Drinking Water Treatment

★ →

Oxidation half-reaction	E_{ox}° , volts
$\frac{1}{2}\text{Br}^- + \frac{1}{2}\text{H}_2\text{O} \rightarrow \frac{1}{2}\text{HOBr} + \frac{1}{2}\text{H}^+ + \text{e}^-$	-1.33
$\frac{1}{2}\text{Mn}^{+2} + \text{H}_2\text{O} \rightarrow \frac{1}{2}\text{MnO}_2(\text{s}) + 2\text{H}^+ + \text{e}^-$	-1.21
$\text{Fe}^{+2} + 3\text{H}_2\text{O} \rightarrow \text{Fe(OH)}_3(\text{s}) + 3\text{H}^+ + \text{e}^-$	-1.01
$\frac{1}{8}\text{NH}_4^+ + \frac{3}{8}\text{H}_2\text{O} \rightarrow \frac{1}{8}\text{NO}_3^- + \frac{1}{4}\text{H}^+ + \text{e}^-$	-0.88
$\frac{1}{2}\text{NO}_2^- + \frac{1}{2}\text{H}_2\text{O} \rightarrow \frac{1}{2}\text{NO}_3^- + \text{H}^+ + \text{e}^-$	-0.84
$\frac{1}{8}\text{H}_2\text{S} + \frac{1}{2}\text{H}_2\text{O} \rightarrow \frac{1}{8}\text{SO}_4^{2-} + \frac{1}{4}\text{H}^+ + \text{e}^-$	-0.30
$\frac{1}{2}\text{H}_2\text{S} \rightarrow \frac{1}{2}\text{S}(\text{s}) + \text{H}^+ + \text{e}^-$	-0.14
$\frac{1}{2}\text{HCOO}^- \rightarrow \frac{1}{2}\text{CO}_2(\text{g}) + \frac{1}{2}\text{H}^+ + \text{e}^-$	+0.29

David Reckhow

CEE 680 #46

7

Oxidation State

- Oxidation state is characterized by an oxidation number
 - the charge one would expect for an atom if it were to dissociate from the surrounding molecule or ion (assigning any shared electrons to the more electronegative atom).
 - may be either a positive or negative number, usually, an integer between -VII and +VII
 - in their elemental forms, e.g. S(s), O₂(aq), atoms have an oxidation number of zero.
 - This concept is useful in balancing chemical equations and performing certain calculations.

David Reckhow

CEE 680 #46

8

Determining oxidation state

- Rule:

- Sum of the oxidation states of all elements in a molecule or ion equals the charge of that molecule or ion

- Conventions:

- H is (+I)
 - Exceptions are H₂, and hydrides
- O is (-II)
 - Exceptions are O₂, and peroxides
- N is (-III) when bound only to C or H
- S is (-II) when bound only to C or H

See Benjamin,
pg. 667

David Reckhow

CEE 680 #45

9

Oxidation States

Stumm & Morgan, 1996;
Table 8.1, pg. 427

Table 8.1. Oxidation State

Rules for Assigning Oxidation States:

- The oxidation state of a monoatomic substance is equal to its electronic charge.
- In a covalent compound, the oxidation state of each atom is the charge remaining on the atom when each shared pair of electrons is assigned completely to the more electronegative of the two atoms sharing them. An electron pair shared by two atoms of the same electronegativity is split between them.
- The sum of oxidation states is equal to zero for molecules, and for ions is equal to the formal charge of the ions.

Examples:

Nitrogen Compounds		Sulfur Compounds		Carbon Compounds	
Substance	Oxidation States	Substance	Oxidation States	Substance	Oxidation States
NH ₄ ⁺	N = -III, H = +I	H ₂ S	S = -II, H = +I	HCO ₃ ⁻	C = +IV
N ₂	N = 0	S ₈ (s)	S = 0	HCOOH	C = +II
NO ₂	N = +III, O = -II	SO ₄ ²⁻	S = +IV, O = -II	C ₆ H ₁₂ O ₆	C = 0
NO ₃ ⁻	N = +V, O = -II	SO ₄ ²⁻	S = +VI, O = -II	CH ₃ OH	C = -II
HCN	N = -III, C = +II, H = +I	S ₂ O ₃ ²⁻	S = +II, O = -II	CH ₄	C = -IV
SCN ⁻	S = -I, C = +III, N = -III	S ₂ O ₆ ²⁻	S = +2.5, O = -II	C ₆ H ₅ COOH	C = -2/7
		S ₂ O ₆ ²⁻	S = +V, O = -II		

David Reckhow

CEE 680 #46

10

Balancing Equations

- The first step in working with oxidation reactions is to identify the role of the reacting species.
 - At least one reactant must be the oxidizing agent (i.e., containing an atom or atoms that become reduced)
 - At least one must be a reducing agent (i.e., containing an atom or atoms that become oxidized).
- The second step is to balance the gain of electrons from the oxidizing agent with the loss of electrons from the reducing agent.
- Next, oxygen atoms are balanced by adding water molecules to one side or another and hydrogens are balanced with H⁺ ions.

David Reckhow

CEE 680 #46

11

Example: Mn & ozone

- As an example consider the oxidation of manganese by ozone .
 - The substance being oxidized is manganese (i.e., the reducing agent), and the one doing the oxidizing (i.e., being itself reduced) is ozone.



- Next the products formed need to be evaluated.
 - It might be known from experience that reduced soluble manganese (i.e., Mn^{+2}) can be oxidized in water to the relatively insoluble manganese dioxide.
 - It might also be known that ozone ultimately forms hydroxide and oxygen after it becomes reduced.

David Reckhow


CEE 680 #46

12

Mn & Ozone (cont.)



- The next step is to determine the oxidation state of all atoms involved.




David Reckhow

CEE 680 #46

13

Mn & Ozone (cont.)

- From this analysis, we conclude:
 - manganese is oxidized from $+II$ to $+IV$, which involves a loss of 2 electrons per atom
 - ozone undergoes a gain of 2 electrons per molecule, as one of the three oxygen atoms goes from an oxidation state of 0 to $-II$.
- The two half-reactions can be written as single electron transfers.
 - These half-reactions are balanced by adding water molecules and H^+ ions to balance oxygen and hydrogen, respectively.




David Reckhow

CEE 680 #46

14

Mn & Ozone (cont.)

- By convention, when hydroxide appears in a half-reaction, additional H⁺ ions are added until all of the hydroxide is converted to water. This is done to the reduction half-reaction.




David Reckhow

CEE 680 #46


15

Mn & Ozone (cont.)

- From this point, it is a simple matter of combining the equations and canceling out terms or portions of terms that appear on both sides.
- At the same time, the standard electrode potentials can be combined to get the overall potential.



$$\begin{aligned} -1.21 \text{ V } (E_{\text{ox}}^{\circ}) \\ + 2.04 \text{ V } (E_{\text{red}}^{\circ}) \end{aligned}$$



$$+ 0.83 \text{ V } (E_{\text{net}}^{\circ})$$

David Reckhow

CEE 680 #46

16

Mn & Ozone (cont.)

- Immediately, it is seen that this reaction will proceed toward the right (the E°_{net} is positive). But how far to the right will it go?
 - To answer this, the previous equation is rearranged to get:

- So for this reaction:

$$K = e^{16.95 E^\circ_{net}}$$

$$K = e^{16.95 \cdot 0.83} = 1.29 \times 10^6$$

David Reckhow

CEE 680 #46

17

Mn & Ozone (cont.)

- and using the concentration quotient from the reaction stoichiometry,

$$\underline{1.29 \times 10^6} = \frac{\{O_{2(aq)}\}^{0.5} \{MnO_{2(s)}\}^{0.5} \{H^+\}}{\{O_{3(aq)}\}^{0.5} \{Mn^{+2}\}^{0.5} \{H_2O\}^{0.5}}$$

- and since the activity of solvents (i.e., water) and solid phases are, by convention, equal to one,

$$\underline{1.29 \times 10^6} = \frac{\{O_{2(aq)}\}^{0.5} \{H^+\}}{\{O_{3(aq)}\}^{0.5} \{Mn^{+2}\}^{0.5}}$$

- Furthermore, if the pH is 7.0 and a dissolved oxygen concentration of 10 mg/L and an ozone concentration of 0.5 mg/L is maintained in the contactor, an equilibrium Mn⁺² concentration of 1.8×10^{-25} M or about 10^{-27} mg/L can be calculated. Thermodynamic principles therefore indicate that this reaction essentially goes to completion.

David Reckhow



CEE 680 #46

18

Mn & Ozone (cont.)

- Now, knowing that the Mn²⁺ should react essentially completely to form manganese dioxide, it might be important to know if ozone can possibly oxidize the manganese dioxide to a higher oxidation state, i.e. permanganate. To examine this, the above ozone equation must first be combined with the reverse of the permanganate reduction equation

+VII



- This allows the net potential to be calculated:

$$E_{\text{net}}^o = E_{\text{ox}}^o + E_{\text{red}}^o = (-1.68V) + (+2.04V) = +0.36V$$

David Reckhow

CEE 680 #46

19

Mn & Ozone (cont.)

- Again, this is a favorable reaction. The equilibrium constant is:

$$\log K = \frac{1}{0.059} E_{\text{net}}^o = \frac{(1)}{0.059} (+0.36V) = 6.1$$

$$K = 10^{6.1}$$

- The equilibrium quotient can now be formulated directly from the balanced equation. Note that neither manganese dioxide (MnO₂) nor water (H₂O) appears in this quotient. This is because both are presumed present at unit activity. Manganese dioxide is a solid and as long as it remains in the system, it is considered to be in a pure, undiluted state. The same may be said for water. As long as the solutes remain dilute, the concentration of water is at its maximum and remains constant.

David Reckhow

CEE 680 #46

20

Mn & Ozone (cont.)

$$K = \frac{\{A_{red}\}^a \{B_{ox}\}^b}{\{A_{ox}\}^a \{B_{red}\}^b}$$

$$K = \frac{[MnO_4]^{0.33} [H^+]^{0.33} [O_2]^{0.5}}{[O_3]^{0.5}} = 10^{6.1}$$

- So under typical conditions where the pH is near neutrality (i.e., $[H^+] = 10^{-7}$), dissolved oxygen is near saturation (i.e., $[O_{2(aq)}] = 3 \times 10^{-4} M$), and the ozone residual is 0.25 mg/L (i.e., $[O_{3(aq)}] = 5 \times 10^{-6} M$), the expected equilibrium permanganate concentration should be:

$$K = \frac{[MnO_4]^{0.33} [10^{-7}]^{0.33} [3 \times 10^{-4}]^{0.5}}{[5 \times 10^{-6}]^{0.5}} = 10^{6.1}$$

David Reckhow

CEE 680 #46

21

Mn & Ozone (cont.)

- and solving for permanganate

$$[MnO_4]^{0.33} = 3.5 \times 10^7$$

$$[MnO_4] = 327 \text{ M}$$

pink water

- Obviously one cannot have 327 moles/liter of permanganate. Nevertheless, this tells us that the system will be forced in this direction so that all of the manganese dioxide would be converted to permanganate. Once the manganese dioxide is gone, the reaction must stop.

David Reckhow

CEE 680 #46

22

p ϵ

- $p\epsilon = -\log\{e^- \}$
- Low $p\epsilon$ means:
 - High $\{e^- \}$
 - Reducing conditions
 - Tend to donate e^-
- $pH = -\log\{H^+ \}$
- Low pH means:
 - High $\{H^+ \}$
 - Acidic conditions
 - Tend to donate H^+

$p\epsilon$ is really a hypothetical construct. There are no free electrons in solution. It is really a measure of the relative tendency to accept or transfer and electron

Equations with $p\epsilon$

- Half reactions as equilibria
 - $Ox + ne^- = Red$

$$K = \frac{[\text{Red}]}{[\text{Ox}] \{e^- \}^n}$$

$$\{e^- \}^n = \frac{1}{K} \frac{[\text{Red}]}{[\text{Ox}]}$$

$$n \log\{e^- \} = -\log K + \log\left(\frac{[\text{Red}]}{[\text{Ox}]}\right)$$

$$p\epsilon = \frac{1}{n} \log K - \frac{1}{n} \log\left(\frac{[\text{Red}]}{[\text{Ox}]}\right)$$

$$p\epsilon = p\epsilon^\circ - \frac{1}{n} \log\left(\frac{[\text{Red}]}{[\text{Ox}]}\right)$$

- [To next lecture](#)