

Updated: 29 April 2020 [Print version](#)

CEE 680: Water Chemistry



Lecture #49
Redox Chemistry: Predominance Diagrams;
spanning the range of p_e
(Stumm & Morgan, Chapt.8)
Benjamin; Chapter 9

David Reckhow CEE 680 #49 1

Chlorine Predominance Diagram

- Features
 - Oxidation state
 - Higher oxidation states (HOCl/OCl^-) in upper region
 - Middle oxidation state (Cl_2) in middle at low pH range
 - Chloride in lower region
 - pH divisions
 - Vertical lines at pK_a values

David Reckhow CEE 680 #48 2



Chlorine Predominance Diagram

- Features
 - Oxidation state
 - Higher oxidation states (HOCl/OCl^-) in upper region
 - Middle oxidation state (Cl_2) in middle at low pH range
 - Chloride in lower region
 - pH divisions
 - Vertical lines at pK_a values

David Reckhow

CEE 680 #49

5

Iron phases

CARBONATE, HALIDE, SULFATE, SULFIDE MINERALS

307

- From Soil Science Literature
 - Pyrite is FeS_2
 - Goethite is FeO(OH)
 - Jarosite is $\text{KFe}^{3+}(\text{OH})_6(\text{SO}_4)_2$






Fig. 6-16. A pH-pe diagram of pyrite, jarosite, amorphous Fe(OH)_3 , goethite, and soluble components at 0.1 MPa total pressure and 25 °C. The activities of solution species are: $(\text{Fe}^{2+}) + (\text{Fe}^{3+}) = 10^{-4}$, $(\text{SO}_4^{2-}) = 5 \times 10^{-3}$, and $(\text{K}^+) = 5 \times 10^{-4}$. Shaded areas represent solid phases. (Diagram after van Breeman, 1982; thermodynamic values for boundaries from Lindsay, 1979).


David Reckhow

CEE 6



Iron (cont.)

- Same as previous, but pH axis limited to 14



From: [Aquatic Chemistry Concepts](#),
by Pankow, 1991 (pg. 536)


David Reckhow

CEE 680 #49

9

Iron (cont.)

- $\text{Fe}_T = 10^{-5} \text{ M}$
- 10^{-3} M carbonate




From: [Aquatic Chemistry Concepts](#), by Pankow, 1991

David Reckhow

CEE 680 #49

10



Fully oxygenated to anaerobic

- pe = 12.6.** In-situ $p_{O_2} = 0.21$ atm. At equilibrium, the dissolved oxygen will be present as O_2 (and a great deal more of H_2O). Since significant dissolved O_2 will be present, the initial pe is fairly high. The carbon, nitrogen, and sulfur will initially be present primarily as HCO_3^- , NO_3^- , and SO_4^{2-} , respectively. $(am)Fe(OH)_{3(s)}$ will be present. The in-situ p_{H_2} will be exceedingly low.
- pe = 12.1.** As the pe is reduced, the first electron acceptor (oxidant) to be reduced in large amounts (but not exclusively) will be the dissolved O_2 , which will be converted to more H_2O . By the time pe = 12.1, the in-situ p_{O_2} will have been reduced to 0.0021 atm.
- pe = 10.0.** When essentially all of the O_2 is exhausted, the next electron acceptor to be reduced in large amounts will be the NO_3^- , which will be converted primarily to N_2 . Once the pe drops to 10, the dominant nitrogen species will be N_2 .
- pe = -2.8.** After most of the NO_3^- has been exhausted, the next electron acceptor to be reduced in large amounts will be the Fe(III). Indeed, once the pe drops to -2.8, $(am)Fe(OH)_{3(s)}$ is no longer present, and we are inside the $FeCO_{3(s)}$ region. Since $Fe_T \ll C_T$, most of the carbon is still present in solution; in solution, we still have $C(IV)_T \simeq 10^{-3} M$.

From: [Aquatic Chemistry Concepts, by Pankow, 1991](#)

5. $pe = -4.9$. After most of the Fe(III) has been exhausted, the next electron acceptor to be reduced in large amounts will be the SO_4^{2-} which will be converted primarily to HS^- . However, since the N_2/NH_4^+ and $\text{HCO}_3^-/\text{CH}_4$ boundaries are only slightly below the $\text{SO}_4^{2-}/\text{HS}^-$ boundary, as SO_4^{2-} begins to accept electrons in significant amounts, N_2 and HCO_3^- will also begin to accept some electrons. Once the pe reaches -4.9 , the dominant sulfur species will be HS^- ; N_2 and HCO_3^- will still be the dominant nitrogen and carbon species, respectively, but significant conversion to NH_4^+ and CH_4 will also have taken place. With C(IV)_T still close to 10^{-3} M , $\text{FeCO}_{3(s)}$ would still be present if it were not for the fact that there was so much sulfur in this system. Indeed, since $\text{FeS}_{(s)}$ is very insoluble, at $pe = -4.9$ the $\text{FeCO}_{3(s)}$ will have been converted to $\text{FeS}_{(s)}$.

6. $pe = -5.3$. Once the pe reaches -5.3 , we drop below the N_2/NH_4^+ line, and NH_4^+ will become the dominant nitrogen species. HCO_3^- will still be the dominant carbon species, but significant amounts of CH_4 will by now also be present.

7. $pe = -6.0$. Once the pe reaches -6.0 , we are below the $\text{HCO}_3^-/\text{CH}_4$ line, and CH_4 is the dominant carbon species. The bulk of the Fe is present as $\text{FeS}_{(s)}$. The level of dissolved Fe(II) is very low and the level of dissolved Fe(III) is exceedingly low.


8. $pe = -8.4$. If the pe continues to be lowered, the last major electron acceptor in the system will be H_2O . Once the pe drops below 8.4 , p_{H_2} will be greater than 1 atm. Since we are still above the line given by Eq. (23.119), $\text{Fe}_{(s)}$ is not present.

From: [Aquatic Chemistry Concepts](#), by Pankow, 1991

David Reckhow

CEE 680 #49


13



Other redox systems

- Phosphorus
- Nitrogen
- Sulfur
- carbon

Stumm & Morgan, 1996;
Fig. 8.9, pg. 461



David Reckhow

CEE 680 #49

15

- To next lecture

DAR

David Reckhow

CEE 680 #49

16