

Updated: 13 November 2013

CEE697K Lecture #16 1

[Print version](#)

CEE 697K

ENVIRONMENTAL REACTION KINETICS

Lecture #16

[Kinetic Modeling](#): Computer Models
[Case Study](#): Chloramination I
Brezonik, pp.

David A. Reckhow Introduction

Kinetic Modeling

2

- Multiple & complex kinetic rate laws are what nature has given us
- Lack of easy analytical solutions
 - ▣ Therefore, need for computer modeling
- Options
 - ▣ Scientist
 - ▣ Mat Lab
 - ▣ [Aquasim](#) (EAWAG)




CEE697K Lecture #16

David A. Reckhow

Chloramines

3

Inorganic chloramines are formed by the reaction of free chlorine with ammonia. The reaction is stepwise, giving monochloramine (equation 1) followed by dichloramine (equation 2). The dichloramine is quite unstable, forming nitrogen gas (equation 3) and some nitrate. This decomposition is responsible for the classic breakpoint chlorination phenomenon.




CEE697K Lecture #16

David A. Reckhow

Chloramines: Breakpoint Curve

4



Chloramination and regulated DBPs

5

- Preponderance of DHAAs; very little THM, TCAA
 - Cowman & Singer, 1996
 - EST 30:1:16
 - Zhang et al., 2000
 - In Natural Organic Matter and Disinfection Byproducts, ACS Symp #761, Barrett, Krasner & Amy, eds.
- HAAs decrease substantially
 - 5-20% of that observed from chlorination (no ammonia addition)
 - Review by Speitel, 1999
- DBP formation increases with decreasing pH; possibly the role of dichloramine
 - Speitel, 1999
 - In Formation and Control of Disinfection Byproducts in Drinking Water, P.C.Singer, ed.
 - Symons et al., 1998
 - AWWARF report

CEE697K Lecture #16

David A. Reckhow

6

- To next lecture

CEE697K Lecture #16

David A. Reckhow