

[Print version](#)

# CEE 697K

## ENVIRONMENTAL REACTION KINETICS

### Lecture #20

[Chloramines revisited](#): bromine, lead and  
bacteria  
Primary Literature

# Model Equations 1 a

Vikesland, P. J., K. Ozekin, et al. (2001).

"[Monochloramine decay in model and distribution system waters.](#)" *Water Research* **35(7)**: 1766-1776.

#	Reaction	Rate coefficient/equilibrium constant (25°C)	References
1	$\text{HOCl} + \text{NH}_3 \rightarrow \text{NH}_2\text{Cl} + \text{H}_2\text{O}$	$k_1 = 1.5 \times 10^{10} \text{ M}^{-1} \text{ h}^{-1}$	<a href="#">Morris and Isaac (1981)</a>
2	$\text{NH}_2\text{Cl} + \text{H}_2\text{O} \rightarrow \text{HOCl} + \text{NH}_3$	$k_2 = 7.6 \times 10^{-2} \text{ h}^{-1}$	<a href="#">Morris and Isaac (1981)</a>
3	$\text{HOCl} + \text{NH}_2\text{Cl} \rightarrow \text{NHCl}_2 + \text{H}_2\text{O}$	$k_3 = 1.0 \times 10^6 \text{ M}^{-1} \text{ h}^{-1}$	<a href="#">Margerum et al. (1978)</a>
4	$\text{NHCl}_2 + \text{H}_2\text{O} \rightarrow \text{HOCl} + \text{NH}_2\text{Cl}$	$k_4 = 2.3 \times 10^{-3} \text{ h}^{-1}$	<a href="#">Margerum et al. (1978)</a>
5	$\text{NH}_2\text{Cl} + \text{NH}_2\text{Cl} \rightarrow \text{NHCl}_2 + \text{NH}_3$	$k_d^a$	Vikesland et al. (2001)
6	$\text{NHCl}_2 + \text{NH}_3 \rightarrow \text{NH}_2\text{Cl} + \text{NH}_2\text{Cl}$	$k_6 = 2.2 \times 10^8 \text{ M}^{-2} \text{ h}^{-1}$	<a href="#">Hand and Margerum (1983)</a>
7	$\text{NHCl}_2 + \text{H}_2\text{O} \rightarrow \text{I}$	$k_7 = 4.0 \times 10^5 \text{ M}^{-1} \text{ h}^{-1}$	<a href="#">Jafvert and Valentine (1987)</a>
8	$\text{I} + \text{NHCl}_2 \rightarrow \text{HOCl} + \text{products}$	$k_8 = 1.0 \times 10^8 \text{ M}^{-1} \text{ h}^{-1}$	<a href="#">Leao (1981)</a>
9	$\text{I} + \text{NH}_2\text{Cl} \rightarrow \text{products}$	$k_9 = 3.0 \times 10^7 \text{ M}^{-1} \text{ h}^{-1}$	<a href="#">Leao (1981)</a>
10	$\text{NH}_2\text{Cl} + \text{NHCl}_2 \rightarrow \text{products}$	$k_{10} = 55.0 \text{ M}^{-1} \text{ h}^{-1}$	<a href="#">Leao (1981)</a>
11	$\text{HOCl} \rightarrow \text{H}^+ + \text{OCl}^-$	$\text{p}K_a = 7.5$	<a href="#">Snoeyink and Jenkins (1980)</a>
12	$\text{NH}_4^+ \rightarrow \text{NH}_3 + \text{H}^+$	$\text{p}K_a = 9.3$	<a href="#">Snoeyink and Jenkins (1980)</a>
13	$\text{H}_2\text{CO}_3 \rightarrow \text{HCO}_3^- + \text{H}^+$	$\text{p}K_a = 6.3$	<a href="#">Snoeyink and Jenkins (1980)</a>
14	$\text{HCO}_3^- \rightarrow \text{CO}_3^{2-} + \text{H}^+$	$\text{p}K_a = 10.3$	<a href="#">Snoeyink and Jenkins (1980)</a>

# What is “I”

3


- NOH first suggested as a key intermediate by Morris, Weil & Culver (1951)?
  - 12th International Congress of Pure and Applied Chemistry: Abstracts of Papers, New York, Sept 10-13, 1951
- Wei (1972) also proposes NOH
  - Wei, Irvine Wen-Tung. "CHLORINE-AMMONIA BREAKPOINT REACTIONS: KINETICS AND MECHANISM." PhD Harvard University; Advisor: J.C. Morris
    - ▣ Acknowledged by Leao (1981)
- Valentine, Jafvert & Leung (1988) say it may or may not be NOH
- Leung & Valentine (1994a, b) say it contains N and Cl
- Maybe it is really several compounds

# Model Equations 1b

#	Reaction	Rate coefficient/equilibrium constant (25°C)	References
1	$\text{HOCl} + \text{NH}_3 \rightarrow \text{NH}_2\text{Cl} + \text{H}_2\text{O}$	$k_1 = 1.5 \times 10^{10} \text{ M}^{-1} \text{ h}^{-1}$	<a href="#">Morris and Isaac (1981)</a>
2	$\text{NH}_2\text{Cl} + \text{H}_2\text{O} \rightarrow \text{HOCl} + \text{NH}_3$	$k_2 = 7.6 \times 10^{-2} \text{ h}^{-1}$	<a href="#">Morris and Isaac (1981)</a>
3	$\text{HOCl} + \text{NH}_2\text{Cl} \rightarrow \text{NHCl}_2 + \text{H}_2\text{O}$	$k_3 = 1.0 \times 10^6 \text{ M}^{-1} \text{ h}^{-1}$	<a href="#">Margerum et al. (1978)</a>
4	$\text{NHCl}_2 + \text{H}_2\text{O} \rightarrow \text{HOCl} + \text{NH}_2\text{Cl}$	$k_4 = 2.3 \times 10^{-3} \text{ h}^{-1}$	<a href="#">Margerum et al. (1978)</a>
5	$\text{NH}_2\text{Cl} + \text{NH}_2\text{Cl} \rightarrow \text{NHCl}_2 + \text{NH}_3$	$k_d^a$	Vikesland et al. (2001)
6	$\text{NHCl}_2 + \text{NH}_3 \rightarrow \text{NH}_2\text{Cl} + \text{NH}_2\text{Cl}$	$k_6 = 2.2 \times 10^8 \text{ M}^{-2} \text{ h}^{-1}$	<a href="#">Hand and Margerum (1983)</a>
7	$\text{NHCl}_2 + \text{OH}^- \rightarrow \text{NOH} + \text{H}^+ + \text{Cl}^-$	$k_7 = 5.5 \times 10^5 \text{ M}^{-1} \text{ h}^{-1}$	<a href="#">Leao &amp; Selleck (1983)</a>
8	$\text{NOH} + \text{NHCl}_2 \rightarrow \text{HOCl} + \text{N}_2 + \text{Cl}^- + \text{H}^+$	$k_8 = 1.0 \times 10^8 \text{ M}^{-1} \text{ h}^{-1}$	<a href="#">Leao (1981)</a>
9	$\text{NOH} + \text{NH}_2\text{Cl} \rightarrow \text{H}_2\text{O} + \text{N}_2 + \text{Cl}^- + \text{H}^+$	$k_9 = 3.0 \times 10^7 \text{ M}^{-1} \text{ h}^{-1}$	<a href="#">Leao (1981)</a>
10	$\text{NH}_2\text{Cl} + \text{NHCl}_2 \rightarrow \text{N}_2 + 3\text{Cl}^- + 3\text{H}^+$	$k_{10} = 55.0 \text{ M}^{-1} \text{ h}^{-1}$	<a href="#">Leao (1981)</a>
11	$\text{HOCl} \rightarrow \text{H}^+ + \text{OCl}^-$	$\text{p}K_a = 7.5$	<a href="#">Snoeyink and Jenkins (1980)</a>
12	$\text{NH}_4^+ \rightarrow \text{NH}_3 + \text{H}^+$	$\text{p}K_a = 9.3$	<a href="#">Snoeyink and Jenkins (1980)</a>
13	$\text{H}_2\text{CO}_3 \rightarrow \text{HCO}_3^- + \text{H}^+$	$\text{p}K_a = 6.3$	<a href="#">Snoeyink and Jenkins (1980)</a>
14	$\text{HCO}_3^- \rightarrow \text{CO}_3^{2-} + \text{H}^+$	$\text{p}K_a = 10.3$	<a href="#">Snoeyink and Jenkins (1980)</a>

# Breakpoint Reactions

5



# Additional Bromide reactions

6

#	Reaction	Rate coefficient/equilibrium constant (25°C)	References
15	$\text{NH}_2\text{Cl} + \text{H}^+ \leftrightarrow \text{NH}_3\text{Cl}^+$	$K = 28 \text{ M}^{-1}$	Gray et al., 1978
15f	$\text{NH}_2\text{Cl} + \text{H}^+ \rightarrow \text{NH}_3\text{Cl}^+$	$k_f = 2.16 \times 10^8 \text{ M}^{-1} \text{ h}^{-1}$	Bousher et al., 1989
15b	$\text{NH}_3\text{Cl}^+ \rightarrow \text{NH}_2\text{Cl} + \text{H}^+$	$k_b = 7.71 \times 10^6 \text{ h}^{-1}$	
16	$\text{NH}_3\text{Cl}^+ + \text{Br}^- \rightarrow \text{NH}_3\text{Br}^+ + \text{Cl}^-$	$k_{\text{Br}} = 1.8 \times 10^8 \text{ M}^{-1} \text{ h}^{-1} \text{a}$	Trofe et al., 1980
17	$\text{NH}_2\text{Cl} + \text{NH}_3\text{Br}^+ \rightarrow \text{NHBrCl} + \text{NH}_4^+$	$k_{\text{fast}}$	Valentine et al., 1998
18	$\text{HOCl} + \text{Br}^- \rightarrow \text{HOBr} + \text{Cl}^-$	$k_{\text{HOCl}} = 4.8 \times 10^6 \text{ M}^{-1} \text{ h}^{-1} \text{b}$	Kumar & Margerum, 1987
19	$\text{HOBr} + \text{NH}_2\text{Cl} \rightarrow \text{NHBrCl} + \text{H}_2\text{O}$	$k_{\text{fast}}$	Valentine et al., 1998
20	$\text{NHBrCl} + \text{H}_2\text{O} \rightarrow \text{NOH} + 2\text{H}^+ + \text{Br}^- + \text{Cl}^-$	$k_{20} = 7.2 \times 10^5 \text{ M}^{-1} \text{ h}^{-1}$	Zhang & Lin, 2013
21	$\text{NOH} + \text{NHBrCl} \rightarrow \text{HOBr} + \text{N}_2 + \text{H}^+ + \text{Cl}^-$	$k_{21} = 5.0 \times 10^8 \text{ M}^{-1} \text{ h}^{-1}$	Zhang & Lin, 2013
22	$\text{NHBrCl} + \text{NH}_2\text{Cl} \rightarrow \text{N}_2 + 3\text{H}^+ + 2\text{Cl}^- + \text{Br}^-$	$k_{\text{fast}}$	Valentine et al., 1998


<sup>a</sup>Values were later adjusted by Zhang & Lin to match results; <sup>b</sup>Reaction#18 only includes neutral reaction

# Breakpoint Reactions

With Bromide;  
Part I – Br analogues

Stable Oxidation Products

7




# Breakpoint Reactions

With Bromide;  
Part II – mixed species

Stable Oxidation  
Products

8




New & faster pathways?




# Chloramines: simplified

9




# Chloramines: with nitroxyl

10




# Chloramines: with chlorine demand

11




# Chloramines: with nitrification

12



# Chloramines: the larger picture

13



# Chloramines: with cometabolism

14





Figure #20

David A. Reckhow

# Chloramines: with pipe reactions

15




# Lead and Nitrite reactions

16

#	Reaction	Rate coefficient/equilibrium constant (25°C)	References
23	$\text{NOH} + \text{PbO}_2 \leftrightarrow \text{Pb}^{+2} + \text{NO}_2^- + \text{OH}^-$	$k_{23} = 1.3 \times 10^5 \text{ m}^{-2}\text{h}^{-1}$	Zhang & Lin, 2013
24	$\text{NO}_2^- + \text{NH}_2\text{Cl} + \text{H}_2\text{O} \rightarrow \text{NO}_3^- + \text{NH}_3 + \text{H}^+ + \text{Cl}^-$	$k_{24} = 4.0 \times 10^7 \text{ M}^{-1}\text{h}^{-1}$	Zhang & Lin, 2013
25	$\text{NO}_2^- + \text{NHCl}_2 + 2\text{H}_2\text{O} \rightarrow \text{NO}_3^- + \text{HOCl} + \text{NH}_3 + \text{H}^+ + \text{Cl}^-$	$k_{25} = 2.0 \times 10^8 \text{ M}^{-1}\text{h}^{-1}$	Zhang & Lin, 2013
26	$\text{NO}_2^- + \text{NHBrCl} + 2\text{H}_2\text{O} \rightarrow \text{NO}_3^- + \text{HOBr} + \text{NH}_3 + \text{H}^+ + \text{Cl}^-$	$k_{26} = 9.0 \times 10^8 \text{ M}^{-1}\text{h}^{-1}$	Zhang & Lin, 2013

Zhang, Y. Y. and Y. P. Lin (2013). "[Release of Pb\(II\) from the reduction of Pb\(IV\) corrosion product PbO<sub>2</sub> induced by bromide-catalyzed monochloramine decomposition.](#)" Environmental Science & Technology **47**: 10931-10938.





Monochloramine cometabolism by *Nitrosomonas europaea* under drinking water conditions

Maestre, J. P., D. G. Wahman, et al. (2013). "[Monochloramine cometabolism by \*Nitrosomonas europaea\* under drinking water conditions.](#)" Water Research **47(13): 4701-4709.**

# Cometabolic Model I

**Table 2 – Process matrix for batch kinetic experiments.**

Kinetic rate	Kinetic rate expression	Reaction stoichiometry				
		$S_{TOTNH_3}$	$S_{NH_2Cl}$	$X_a$	$S_{NO_2}$	$S_{UAP}$
Ammonia first-order	$k_{1TOTNH_3} X_a S_{TOTNH_3} \alpha_1$	-1			1	$f_{UAP}$
Ammonia monod	$\frac{k_{TOTNH_3} X_a S_{TOTNH_3} \alpha_1}{K_{S_{NH_3-N}} + S_{TOTNH_3} \alpha_1}$	-1			1	$f_{UAP}$
First-order cometabolism	$k_{1NH_2Cl} X_a S_{NH_2Cl}$		-1		1	
First-order-reductant cometabolism	$k_{1NH_2Cl} X_a S_{NH_2Cl} \left( \frac{S_{TOTNH_3} \alpha_1}{K_{S_{NH_3-N}} + S_{TOTNH_3} \alpha_1} \right)$		-1		1	
Biomass reactivity	$k_{Biomass} X S_{NH_2Cl}$	1	-1			
Biomass inactivation	$k_{Inact} X_a S_{NH_2Cl}$			-1		
UAP Reactivity	$k_{UAP} S_{UAP} S_{NH_2Cl}$	1	-1			-1

$f_{UAP}$  – UAP formation fraction from TOTNH<sub>3</sub> degradation, moles UAP formed/moles TOTNH<sub>3</sub> degraded.

$k_{1TOTNH_3}$  – ammonia first-order rate constant, moles TOTNH<sub>3</sub>-L/(moles NH<sub>3</sub>-N mg TSS day).

$X_a$  – active biomass concentration, mg TSS/L.

$S_{TOTNH_3}$  – TOTNH<sub>3</sub> concentration, moles TOTNH<sub>3</sub>/L.

$\alpha_1$  – NH<sub>3</sub>-N fraction of TOTNH<sub>3</sub>.

$k_{TOTNH_3}$  – ammonia maximum specific rate of degradation, moles TOTNH<sub>3</sub>/mg TSS-day.

$K_{S_{NH_3-N}}$  – ammonia half-saturation constant, moles NH<sub>3</sub>-N/L.

$k_{1NH_2Cl}$  – monochloramine first-order cometabolism rate constant, L/mg TSS-day.

$S_{NH_2Cl}$  – monochloramine concentration, moles Cl<sub>2</sub>/L.

$k_{Biomass}$  – monochloramine reaction with biomass rate constant, L/mg TSS-day.

$X$  – initial biomass concentration, mg TSS/L.

$k_{Inact}$  – active biomass inactivation rate constant, L/moles Cl<sub>2</sub>-day.

$k_{UAP}$  – monochloramine reaction rate constant with UAP, L/moles UAP-day.

$S_{UAP}$  – UAP concentration, moles UAP/L.

Maestre, J. P., D. G. Wahman, et al. (2013). "Monochloramine cometabolism by *Nitrosomonas europaea* under drinking water conditions." [Water Research 47\(13\): 4701-4709.](#)

# Cometabolic Model II

**Table 3 – Summary and comparison of kinetic parameters for *N. europaea*.**

Parameter	Description	Units	Estimate	Standard deviation
$k_{TOTNH_3}$	Ammonia maximum specific rate of degradation	mg TOTNH <sub>3</sub> /mg TSS–day	2.9	Fixed constant
$k_{S_{NH_3-N}}$	Ammonia half-saturation constant	mg NH <sub>3</sub> –N/L	0.13	0.0035
$k_{1TOTNH_3}$	Ammonia first-order rate constant	$\frac{(\text{mg TOTNH}_3)(L)}{(\text{mg NH}_3 - N)(\text{mg TSS})(\text{day})}$	22.3	0.6
$k_{1TOTNH_3}^* \alpha_{1_{pH8.3}}$	Ammonia first-order rate constant at pH 8.3	L/mg TSS–day	2.3	0.06
$k_{Biomass}$	Monochloramine reaction with biomass rate constant	L/mg TSS–day	0.18	0.031
$k_{Inact}$	Active biomass inactivation rate constant	L/mg Cl <sub>2</sub> –day	224	22
$f_{UAP}$	UAP formation fraction from TOTNH <sub>3</sub> metabolism	mole UAP/mole TOTNH <sub>3</sub>	0.029	0.0091
$k_{UAP}$	Monochloramine reaction rate constant with UAP	1/M–day – L/moles UAP–day	$1.85 \times 10^7$	$1.54 \times 10^7$
$k_{1NH_2Cl}$	Monochloramine first-order cometabolism rate constant	L/mg TSS–day	2.1	0.53
$k_{1TCM}^a$	Chloroform first-order cometabolism rate constant	L/mg TSS–day	0.10	
$k_{1TBM}^a$	Bromoform first-order cometabolism rate constant	L/mg TSS–day	0.23	

a Chloroform and bromoform rate constants from Wahman et al. (2005).

Maestre, J. P., D. G. Wahman, et al. (2013). "[Monochloramine cometabolism by Nitrosomonas europaea under drinking water conditions.](#)" [Water Research](#) **47(13): 4701-4709.**




Fig. 1. Comparison of total free ammonia (TOTNH<sub>3</sub>) model simulations and experimental data for a subset of positive control experiments shown over a range of initial TOTNH<sub>3</sub> concentrations. Initial conditions: pH 8.3 and 4 mM bicarbonate buffer.

Maestre, J. P., D. G. Wahman, et al. (2013). "[Monochloramine cometabolism by Nitrosomonas europaea under drinking water conditions.](#)" [Water Research 47\(13\): 4701-4709.](#)




Fig. 2. Comparison of monochloramine (NH<sub>2</sub>Cl) and total free ammonia (TOTNH<sub>3</sub>) model simulations and experimental data in Experiment J.

Comet. = Cometabolism and Neg. Cont. = Negative Control. Initial conditions: pH 8.3, 4 mM bicarbonate buffer, and 5:1 Cl<sub>2</sub>:N mass ratio.

Maestre, J. P., D. G. Wahman, et al. (2013). "[Monochloramine cometabolism by Nitrosomonas europaea under drinking water conditions.](#)" [Water Research](#) **47(13): 4701-4709.**




Fig. 3. Comparison of monochloramine ( $\text{NH}_2\text{Cl}$ ) model simulations and experimental data for a subset of negative control experiments for a range of initial  $\text{NH}_2\text{Cl}$  concentrations. Experiments detail the reaction of  $\text{NH}_2\text{Cl}$  with chlorobenzene inactivated biomass. Initial conditions: pH 8.3 and 4 mM bicarbonate buffer.

Maestre, J. P., D. G. Wahman, et al. (2013). "[Monochloramine cometabolism by \*Nitrosomonas europaea\* under drinking water conditions.](#)" [Water Research](#) **47(13): 4701-4709.**




Figure 1 Distribution of inorganic chloramines depending on (A) molar ratio N/Cl, and on (B) pH, according to Cimetière [\[5\]](#) .

Said Kinani , Bertille Richard , Yasmine Souissi , Stéphane Bouchonnet

**Analysis of inorganic chloramines in water**

TrAC Trends in Analytical Chemistry Volume 33 2012 55 - 67

<http://dx.doi.org/10.1016/j.trac.2011.10.006>

# Vikesland modification

24

□ Modified version of equation #5 for carbonate

$$\square k_d = k_H^+ [H^+] + k_{H_2CO_3} [H_2CO_3] + k_{HCO_3} [HCO_3]$$

□ Where

■  $k_H^+ = 2.5 \times 10^7 \text{ M}^{-2}\text{h}^{-1}$

■  $k_{H_2CO_3} = 4 \times 10^4 \text{ M}^{-2}\text{h}^{-1}$

■  $k_{HCO_3} = 800 \text{ M}^{-2}\text{h}^{-1}$

■ I is the unidentified monochloramine auto-decomposition intermediate

Vikesland, P. J., K. Ozekin, et al. (2001).

"Monochloramine decay in model and distribution system waters." *Water Research* **35(7): 1766-1776.**






# Temperature Effects

25

Reaction	Rate coefficient/equilibrium constant	References
$\text{HOCl} + \text{NH}_3 \rightarrow \text{NH}_2\text{Cl} + \text{H}_2\text{O}$	$k_{1,1} = 2.37 \times 10^{12} \exp(-1510/T) \text{ M}^{-1} \text{ h}^{-1}$	<a href="#">Morris and Isaac (1981)</a>
$\text{NH}_2\text{Cl} + \text{H}_2\text{O} \rightarrow \text{HOCl} + \text{NH}_3$	$k_{1,2} = 6.7 \times 10^{11} \exp(-8800/T) \text{ h}^{-1}$	<a href="#">Morris and Isaac (1981)</a>
$\text{HOCl} + \text{NH}_2\text{Cl} \rightarrow \text{NHCl}_2 + \text{H}_2\text{O}$	$k_{1,3} = 1.08 \times 10^9 \exp(-2010/T) \text{ M}^{-1} \text{ h}^{-1}$	<a href="#">Margerum et al. (1978)</a>
$\text{NH}_2\text{Cl} + \text{NH}_2\text{Cl} \rightarrow \text{NHCl}_2 + \text{NH}_3$	$k_{1,5} = k_{\text{H}^+}[\text{H}^+] + k_{\text{HCO}_3^-}[\text{HCO}_3^-] + k_{\text{H}_2\text{CO}_3}[\text{H}_2\text{CO}_3]$	
	$k_{\text{H}^+} = 3.78 \times 10^{10} \exp(-2169/T) \text{ M}^{-2} \text{ h}^{-1}$	<a href="#">Granstrom (1954)</a>
	$k_{\text{HCO}_3^-} = 1.5 \times 10^{35} \exp(-22144/T) \text{ M}^{-2} \text{ h}^{-1}$	Vikesland et al. (2001)
	$k_{\text{H}_2\text{CO}_3} = 2.95 \times 10^{10} \exp(-4026/T) \text{ M}^{-2} \text{ h}^{-1}$	Vikesland et al. (2001)
$\text{H}_2\text{CO}_3 \rightleftharpoons \text{HCO}_3^- + \text{H}^+$	$\text{p}k_a = 1.48 \times 10^{-4} (T) - 9.39 \times 10^{-2} (T) + 21.2$	<a href="#">Snoeyink and Jenkins (1980)</a>
$\text{HCO}_3^- \rightleftharpoons \text{H}^+ + \text{CO}_3^{2-}$	$\text{p}k_a = 1.19 \times 10^{-4} (T) - 7.99 \times 10^{-2} (T) + 23.6$	<a href="#">Snoeyink and Jenkins (1980)</a>
$\text{NH}_4^+ \rightleftharpoons \text{NH}_3 + \text{H}^+$	$\text{p}k_a = 1.03 \times 10^{-4} (T) - 9.21 \times 10^{-2} (T) + 27.6$	<a href="#">Bates and Pinching (1950)</a>
$\text{HOCl} \rightleftharpoons \text{OCl}^- + \text{H}^+$	$\text{p}k_a = 1.18 \times 10^{-4} (T) - 7.86 \times 10^{-2} (T) + 20.5$	

## Br<sup>-</sup>-catalyzed NH<sub>2</sub>Cl decomposition & PbO<sub>2</sub> reduction



Zhang, Y. Y. and Y. P. Lin (2013). "[Release of Pb\(II\) from the reduction of Pb\(IV\) corrosion product PbO<sub>2</sub> induced by bromide-catalyzed monochloramine decomposition.](#)" *Environmental Science & Technology* **47**: **10931-10938**.

- To next lecture