

[Print version](#)

## CEE 697z Organic Compounds in Water and Wastewater

PPCPs: Source Loads and Observations in  
Natural Systems

Lecture #18


For Background see:

<http://www.ecs.umass.edu/eve/background/chemicals/PPCPs/PPCP%20occurrence.html>

|

CEE 697z - Lecture #18 David A. Reckhow

### EDC/PPCP Sources



## Estimating Source Terms

- ▶ Use-based calculations (e.g., Sedlak)
  - ▶ Get national or regional use data
  - ▶ Estimate non-metabolized/adsorbed fraction
  - ▶ Estimate removal across conventional WWT
- ▶ Real WW effluent monitoring
  - ▶ Highly variable based on date, time, location, processes, climate, etc

▶ 3

CEE 697z - Lecture #18 David A. Reckhow

## Sui et al., 2011

**ENVIRONMENTAL**  
Science & Technology

ARTICLE

pubs.acs.org/est


### Seasonal Variation in the Occurrence and Removal of Pharmaceuticals and Personal Care Products in Different Biological Wastewater Treatment Processes

Qian Sui, Jun Huang, Shubo Deng, Weiwei Chen, and Gang Yu\*

School of Environment, THU – VEOLIA Joint Research Center for Advanced Environmental Technology, Tsinghua University, Beijing 100084, China

Supporting Information

**ABSTRACT:** The occurrence of 12 pharmaceuticals and personal care products (PPCPs) in two wastewater treatment plants in Beijing was studied monthly over the course of one year. The removal of PPCPs by three biological treatment processes including conventional activated sludge (CAS), biological nutrient removal (BNR), and membrane bioreactor (MBR) was compared during different seasons. Seasonal variations of PPCPs in the wastewater influent were discrepant, while in the wastewater effluent, most PPCPs had lower concentrations in the summer than in the winter. For the easily biodegradable PPCPs, the performance of MBR was demonstrated to be more stable than CAS or BNR especially during winter months. Diclofenac, trimethoprim, metoprolol, and gemfibrozil could be moderately removed by MBR, while their removal by CAS and BNR was much lower or even negligible. Nevertheless, no removal was achieved regardless of the season or the treatment processes for the recalcitrant PPCPs. Studies on the contribution of each tank of the MBR process to the total removal of four biodegradable PPCPs indicated theoxic tank was the most important unit, whereas membrane filtration made a negligible contribution to their elimination.



Different wastewater treatment processes in Beijing

Sui, Q., Huang, J., Deng, S.B., Chen, W.W. and Yu, G. (2011) [Seasonal Variation in the Occurrence and Removal of Pharmaceuticals and Personal Care Products in Different Biological Wastewater Treatment Processes](#).

Environmental Science & Technology 45(8), 3341-3348.

A. Reckhow

## Compounds Studied

- ▶ **BF** → Bezafibrate
- ▶ **CBZ** → Carbamazepine
- ▶ **CF** → Caffeine
- ▶ **CP** → Chloramphenicol
- ▶ **DEET** → N,N-diethyl-m-toluamide
- ▶ **DF** → Diclofenac
- ▶ **GF** → Gemfibrozil
- ▶ **MTP** → Metoprolol
- ▶ **SP** → Sulpiride
- ▶ **TP** → Trimethoprim

▶ 5


CEE 697z - Lecture #18 David A. Reckhow

## Seasonal Variability


Seasonal variation in the concentrations of some PPCPs in the wastewater influents (a-d) and effluents (e-h). The symbols represent the mean concentration, and error bars represent the maximum and minimum concentration in CAS, MBR, and BNR processes.



Published in: Qian Sui, Jun Huang, Shubo Deng, Weiwei Chen, Gang Yu, *Environ. Sci. Technol.* 2011, 45, 3341-3348.



## Process Removal Efficiencies



Comparison of the overall removal efficiencies by:

- Conventional Activated Sludge (CAS)
- Biological Nutrient Removal (BNR), and
- Membrane Bioreactor (MBR) processes.




Published in: Qian Sui, Jun Huang, Shubo Deng, Weiwei Chen, Gang Yu, *Environ. Sci. Technol.* 2011, 45, 3341-3348  
 DOI: 10.1021/es200248d  
 Copyright © 2011 American Chemical Society

CEE 677z - Lecture #18 David A. Reckhow


## Seasonal variability for removal

Seasonal variation in the removal efficiencies of PPCPs during the whole year: comparison among MBR and other two biological treatment processes.




Published in: Qian Sui, Jun Huang, Shubo Deng, Weiwei Chen, Gang Yu, *Environ. Sci. Technol.* 2011, 45, 3341-3348  
 DOI: 10.1021/es200248d  
 Copyright © 2011 American Chemical Society

CEE 677z - Lecture #18 David A. Reckhow



## Generic Mechanistic View



**Figure 1** | Micro-pollutant (MP) fractions and processes, influencing MP removal in wastewater. Compiled based on studies by Critticle (1993); Alvarez-Cohen & Speitel (2001); Torres & Joss (2006); Melcer et al. (2007); Monteith et al. (2008); Lindblom et al. (2009); Barret et al. (2010a); Płósz et al. (2010b,c). DCM, dissolved and colloidal matter.

Płósz, B.G., Benedetti, L., Daigger, G.T., Langford, K.H., Larsen, H.F., Monteith, H., Ort, C., Seth, R., Steyer, J.P. and Vanrolleghem, P.A. (2013) Modelling micro-pollutant fate in wastewater collection and treatment systems: status and challenges. *Water Science and Technology* 67(1), 1-15.

Nelson et al., 2010

**ENVIRONMENTAL**  
 Science & Technology

ARTICLE

pubs.acs.org/est

### Diurnal Variability of Pharmaceutical, Personal Care Product, Estrogen and Alkylphenol Concentrations in Effluent from a Tertiary Wastewater Treatment Facility

Eric D. Nelson,\* Huy Do, Roger S. Lewis, and Steve A. Carr

San Jose Creek Water Quality Laboratory, Sanitation Districts of Los Angeles County, 1965 Workman Mill Rd., Whittier California 90601, United States

Supporting Information


**ABSTRACT:** Hourly samples of tertiary wastewater effluent were analyzed for 30 pharmaceuticals, personal care products, estrogenic steroids, and alkylphenols in order to better understand the rate at which these compounds enter the environment. Several distinct patterns of daily cycling were observed, and were characterized as three separate categories. The concentrations of compounds such as trimethoprim, sulfamethoxazole, naproxen, estrone, and triclosan varied greatly during a daily cycle, with relative standard deviations exceeding 100% of their daily mean. Less extreme daily cycles were seen for other compounds such as azithromycin, atenolol, *tert*-octylphenol, iopromide and gemfibrozil. Peak concentrations for most compounds occurred in the early evening (5–8 pm). However, some compounds including carbamazepine, primidone, fluoxetine, and triclocarban exhibited little or no variability.

Nelson, E.D., Do, H., Lewis, R.S. and Carr, S.A. (2010) [Diurnal Variability of Pharmaceutical, Personal Care Product, Estrogen and Alkylphenol Concentrations in Effluent from a Tertiary Wastewater Treatment Facility](#) *Environmental Science & Technology* 45(4), 1228-1234.

▶ 11

CEE 697z - Lecture #18 David A. Reckhow


## Tertiary WRF, Los Angeles

 ACS Publications  
 High quality. High impact.


Simplified schematic of treatment plant train, and location of sample collection.

Published in: Eric D. Nelson; Huy Do; Roger S. Lewis; Steve A. Carr; *Environ. Sci. Technol.* 2011, 45, 1228-1234.  
 DOI: 10.1021/es102452f  
 Copyright © 2010 American Chemical Society

## Flow Variability



Composite daily plant flow (combined of three events). Less consistent effluent flow is due to periodic back flushing of 3<sup>o</sup> filter beds.


Published in: Eric D. Nelson; Huy Do; Roger S. Lewis; Steve A. Carr; *Environ. Sci. Technol.* 2011, 45, 1228-1234.  
DOI: 10.1021/es102452f  
Copyright © 2010 American Chemical Society

CEE 677z - Lecture #18 David A. Reckhow

## Diurnal Variability in WW

Hourly concentrations of selected analytes in wastewater effluent.

Closed circles represent samples from July 2008, open circles represent May 2009, and open triangles represent October 2009.



Published in: Eric D. Nelson; Huy Do; Roger S. Lewis; Steve A. Carr; *Environ. Sci. Technol.* 2011, 45, 1228-1234.

► [To next lecture](#)

