

Sorption of PPCPs

Organic compounds in water and wastewater

Soonmi Kim

Outline

- ◆ Introduction
- ◆ Studies; sorption of PPCPs


Introduction

Sorption?

Sorption is a physical and chemical process by which one substance becomes attached to another

Adsorption




Accumulation of the molecular species at the surface rather than in the bulk of the solid or liquid





Absorption

Assimilation of molecular species throughout the bulk of the solid or liquid


Introduction



Introduction



Wastewater treatment systems




Factors affecting Sorption

1) Sorbent properties



2) Sorbate properties



3) Solution properties




Factors affecting Sorption

1) Sorbents properties



2) Sorbate properties



3) Solution properties



Sorbents

I. Types of sorbent

1) Biosolids (Sludge)



- Primary sludge
- Secondary sludge

2) Soils

- Sand
- Silt
- Clay

3) Coagulants

- Al, Fe, or Mn complex





2. Surface area of sorbents

Factors affecting Sorption

1) Sorbent properties



2) Sorbate properties



3) Solution properties



Sorbate properties

Molecular Weight (MW)

- in general, as increase MW, solubility decreases, adsorption increases
- as increase MW, diffusivity decreases, so rate of transport decreases
- at very high MW, may exclude some compounds from small pores

Polarity


- in general, if more polar, more hydrophilic, less absorption (stronger interaction with polar H_2O)
- as more non-polar, more hydrophobic, more absorption (i.e. increase in aliphatic carbon chain length for carboxylic acids results in increased absorption)
- number and type of substituent atoms or groups on organic carbon structure affects extent of absorption

Factors affecting Sorption

1) Sorbents properties



2) Sorbate properties



3) Solution properties



Solution properties (cont.)

pH

- affects ionization of sorbate and surface of sorbent
- in general, as pH decreases, sorption increases
- uncharged sorbates adsorb more than charged (ionized) sorbates

Ionic Strength

- in general, more sorption as ionic strength increases
- better adsorption of DOC as increase Ca^{+2} , Na^{+}

Temperature

- extent of sorption increases as temperature decreases
- rate of sorption decreases as temperature decreases

Solution properties

Competing sorbates









- sorbate with stronger affinity (binding) for surface can displace sorbate with weaker affinity
- most real waters involve mixtures of sorbates
- changes in mixture can affect sorption
- for a heterogeneous sorbate measured by a composite parameter such as TOC (or COD), extent of sorption can depend on initial concentration used in batch experiment



Presence/absence of dissolved oxygen

- sorption capacity increased in presence of $O_2(aq)$

Possible interactions

Table 1. Characteristic interactions associated with categories of adsorption

Category and Characteristic Interaction	Representation of Interaction	Interaction Range
CHEMICAL		
Covalent		Short Range
H-bonding [Fotyma, Mercik 1992]		
Hydrogen Bond		Short Range
ELECTROSTATIC		
Ion-Ion		$1/r$
Ion-Dipole		$1/r^2$
PHYSICAL		
Dipole-Dipole (Coulombic)		$1/r^3$
(Keesom energy)		$1/r^6$
Dipole-Induced Dipole (Debye energy)		$1/r^6$
Instantaneous Dipole-Induced Dipole (London dispersion energy)		$1/r^6$



*Adapted from Israelachvili, 1985.

Outline

- ◆ Introduction
- ◆ Studies; sorption of PPCPs

Sorption coefficient (K_d)

- The ratio of the concentrations of a compound in the sorbent phase and in the solution phase at equilibrium; Linear isotherm

$$K_d = C_{\text{sorbed}} / C_{\text{dissolved}}$$

$$q = k_{\text{lin}} c$$

C_{sorbed} : concentration of solbate in sorbent ($\mu\text{g/kg}$)

$C_{\text{dissolved}}$: concentration of solbate in solution ($\mu\text{g/L}$)

q = Mass adsorbed / Mass of adsorbent


Other isotherms

Freundlich

$$q = k_f c^n$$


Langmuir

$$q = q_{\text{max}} \frac{K_L c}{1 + K_L c}$$




Standard methods for K_d and K_F determination

I) Column displacement studies



2) Batch sorption experiments



24hr



Factors affecting determination of sorption coefficient

1. Phases associated sorption

- Solid phase
- PPCPs in aqueous phase
- DOM (Dissolved organic Matter) in aqueous phase

2. PPCPs properties

- Hydrophobic PPCPs
- Non-hydrophobic ionisable PPCPs

K_d in soils




Table 4. Sorption coefficients for gemfibrozil to sand, a sandy loam, and a silt loam soil

Sorbent	Freundlich equation			Linear equation		
	K_f	$1/n$	r^2	K_d	r^2	Log K_{oc}
Sand	0.24	0.68	0.96	0.12	0.94	2.08
Sandy loam	1.31	0.83	1.00	1.06	1.00	1.91
Silt loam	12.38	1.37	0.97	9.2	0.99	2.57

Yu Fang (2011)

Table 1 A summary table of PPCP sorption on soils or sediments

Solute chemical names	pK _a	S _w (mg/L)	logK _{ow}	MW	Sorbent description	pH	TOC (%)	K _d (mL/g)	Reference
Bisphenol A (BPA)	9.59–10.2	120– 300	3.4	228.3	Sediment	7.15–7.71	2.06–6.29	3.7–11.5 ^a	Zeng et al. 2006
					Soil	8.9	1.9	2.75 ^a	Ying et al. 2003
					Soil	6.9–8.13	0.92–2.11	4.94–8.62 ^a	Fent et al. 2003
					Minerals	4–10		12.3–212.8 ^a	Shareef et al. 2006
					Zeolite			11.2 ^a	Tsai et al. 2006a
17β-Estradiol (E2)	N/A	3.6	3.1 to 4.0	272.3	Soil	7.2–7.6	0.8–1.4	16.0–29.6 ^a	Stumpe and Marschner 2007
					Soil		3.3–9.2	80.2–455.5 ^a	Casey et al. 2003
					Soil	7.9	9.2	84.41	Casey et al. 2005
					Soil			3.6–83.2	Lee et al. 2003
					Minerals			4.9–177.2	Van Emmerik et al. 2003
17α-Ethinyl Estradiol (EE2)	10.4	11–44	4.15	296.4	Minerals		0	11.3–40 ^a	Casey et al. 2003
					Sludge	7.1	27.7	584	Andersen et al. 2005
					Soil		0.95–1.88	53.7–97.7 ^a	Yu and Huang 2005
					Soil	8.9	1.9	6.98 ^a	Ying et al. 2003
					Soil	6.4–7.3	0.22–2.91	2.33–23.4	Lee et al. 2003
Carbadox (CBX)	N/A	N/A	–1.40 to 0.15	262	Minerals	4–10		41.4–744.9 ^a	Shareef et al. 2006
					Soil	4.4–7.3	0.22–2.39	1.4–154	Strock et al. 2005
Desoxycarbadox (DCBX)	N/A	N/A	1.9 to 2.1	230	Kaolinite	4.63		19.8	Strock et al. 2005
					Soil	4.3–6.9	0.36–2.39	7.07–62.4 ^a	Strock et al. 2005
Tetracycline (TC)	3.3; 7.7; 9.7	231– 52000	–1.97 to – 0.47	444.4	Clay	4.67	0	7.6	Strock et al. 2005
					Soil	3.8–7.5	0.48–2.91	3102– 312447	Sassman and Lee 2005
					HS	7		1.0–5.7E4	Pils and Laird 2007
					Al or Fe oxide	5.3		41.8–133 ^a	Gu and Karthikeyan 2005a
					MMLT	5.5		865 ^a	Figuerola et al. 2004
Oxytetracycline (OTC)	3.27; 7.32; 9.11	300	–1.97 to – 0.45	460	Clay	7		2.2–6.5E4	Pils and Laird 2007
					Soil	3.8–7.5	0.48–2.91	1229– 269097	Sassman and Lee 2005
								950–7200	ter Laak et al. 2006a
					MMLT	1.5–11		2691– 33884 ^a	Kulshrestha et al. 2004
Sulfathiazole (STZ)	2.4; 7.1	590	1	255.3	MMLT	4–7.5		0.5–1.5	Kahle and Stamm 2007
					Ferrihydrite	5.3–7.2		3–20	Kahle and Stamm 2007
Sulfapyridine (SPY)	8.4	270	0.35	249.3	Soil			0.4–3.5	ter Laak et al. 2006a
					Whole soil	7	1.61	1.75 ^a	Thiele-Bruhn et al. 2004
Sulfamethazine (SMZ)	2.3; 7.4	1500	0.8	278.33	Mineral			4	Gu and Karthikeyan 2005b
					Whole soil				
p-Aminobenzoic acid (pABA)	4.9	6100	0.83	137.1	Soil	7.2–7.5	0.94–1.8	2.4–4.2 ^a	Accinelli et al. 2007
					Mineral			7.8–14.2 ^a	Gao and Pedersen 2005
					Soil	7	1.61	0.47 ^a	Thiele-Bruhn et al. 2004
					Coarse silt	6.4	0.1	1.82 ^a	Thiele-Bruhn et al. 2004

S_w solubility in water, MW molecular weight, N/A not available, MMLT montmorillonite

^a K_d was calculated at 10 µg/L if the data was provided as fitting results using the Freundlich or Langmuir model.

Sorption between PPCPs (VPs) and DOM




FIGURE 3. Plot of the $\log K_{d,DOM}$ data against hydrophobicity expressed as $\log K_{ow}$. The solid line is a regression line obtained for a wide range of neutral organic chemicals (32).

Johannes Tolls(2001)

TABLE 3. Data on Sorption of VPs to DOM^a

compound	corollary information	K_d (L/kg)	K_{oc} (L/kg)	ref
tetracycline	AHA, pH adjusted to 4.55, ED	2 060	6 059	24
	pH adjusted to 6.14, ED	1 430	4 206	24
oxolinic acid	AHA, pH range from 3 to 8, SPME	935–8 350	7 400–31 600	25
flumequine	AHA, pH range from 3 to 8, SPME	2 516–10 750	2 750–24 500	25
sarafloxacin	AHA, pH range from 3 to 8, SPME	18 700–52 700	55 000–155 000	25
enrofloxacin	HAS, pH 9.2, EM ^b	110		26
ciprofloxacin	HAS, pH 9.2, EM ^b	250		26
norfloxacin	HAS, pH 9.2, EM ^b	500		26
danofloxacin	HAS, pH 9.2, EM ^b	630		26
ofloxacin	HAS, pH 9.2, EM ^b	100		26
enro-CO ₂	HAS, pH 9.2, EM ^b	250		26

^a AHA and HAS stand for Aldrich humic acid and for humic acid from a soil, respectively. The experimental techniques employed were equilibrium dialysis (ED), solid-phase microextraction (SPME), and electrophoretic mobility. K_d and K_{oc} are the sorption and organic carbon-normalized sorption coefficients, respectively. ^b Organic carbon content of humic acid is not reported.

Hydrophobicity (Cont.)

- Karickhoff (1979) reported that K_d of environmental chemicals are strongly correlated to the organic carbon content ($f_{oc}(\%)$)
- During(2002), Bowman(2002), Holthaus(2002), Loffredo(2006), Maskaoui(2007), and Uslu(2008) reported that both Freundlich sorption parameter K_F and Linear sorption parameter K_d of PPCPs are found to be positively related to organic carbon content
- Most of PPCPs are moderately hydrophobic compounds (Bo Pan;2009)

So,

$$K_{OC} = \frac{K_d}{f_{OC}}$$

K_{oc} : organic carbon partition coefficient (L/kg)

K_d : linear sorption coefficient (L/kg)

f_{oc} : fraction of organic carbon (%)

Hydrophobicity (Cont.)








- Franco(2008) and Karickhoff (1979) reported the K_{oc} is correlated to the K_{ow} (Octanol-water distribution coefficient)
 - $\text{Log } K_{ow} = \text{positive}$: Hydrophobic
 - $\text{Log } K_{ow} = \text{negative}$: Non-hydrophobic
- Sablijic (1995) found that the K_{oc} for non-hydrophobic and ionisable compounds is pH-dependent
- And, Sablijic (1995) determined the correlation between the K_{oc} and the K_{ow} depending on physical and chemical properties of organic compounds

Table 5. List of all derived QSAR models for soil sorption with their chemical domains.

Model #	Regression Equation	Chemical Domain
1	$\log K_{oc} = 0.70 + 0.52 \cdot \log K_{ow}$	Predominantly hydrophobics
2	$\log K_{oc} = 0.10 + 0.81 \cdot \log K_{ow}$	Predominantly hydrophobics
3	$\log K_{oc} = 1.02 + 0.52 \cdot \log K_{ow}$	Nonhydrophobics
4	$\log K_{oc} = 0.90 + 0.63 \cdot \log K_{ow}$	Phenols, anilines, benzonitriles, & nitrobenzenes
5	$\log K_{oc} = 1.09 + 0.47 \cdot \log K_{ow}$	Acetanilides, carbamates, esters, phenylureas, phosphates, triazines, triazoles, & uracils
6	$\log K_{oc} = 0.50 + 0.47 \cdot \log K_{ow}$	Alcohols & organic acids
7	$\log K_{oc} = 1.12 + 0.40 \cdot \log K_{ow}$	Acetanilides
8	$\log K_{oc} = 0.50 + 0.39 \cdot \log K_{ow}$	Alcohols
9	$\log K_{oc} = 1.25 + 0.33 \cdot \log K_{ow}$	Amides
10	$\log K_{oc} = 0.85 + 0.62 \cdot \log K_{ow}$	Anilines
11	$\log K_{oc} = 1.14 + 0.365 \cdot \log K_{ow}$	Carbamates
12	$\log K_{oc} = 1.92 + 0.38 \cdot \log K_{ow}$	Dinitroanilines
13	$\log K_{oc} = 1.05 + 0.49 \cdot \log K_{ow}$	Esters
14	$\log K_{oc} = 0.55 + 0.77 \cdot \log K_{ow}$	Nitrobenzenes
15	$\log K_{oc} = 0.32 + 0.60 \cdot \log K_{ow}$	Organic acids
16	$\log K_{oc} = 1.08 + 0.57 \cdot \log K_{ow}$	Phenols & benzonitriles
17	$\log K_{oc} = 1.05 + 0.49 \cdot \log K_{ow}$	Phenylureas
18	$\log K_{oc} = 1.17 + 0.49 \cdot \log K_{ow}$	Phosphates
19	$\log K_{oc} = 1.50 + 0.30 \cdot \log K_{ow}$	Triazines
20	$\log K_{oc} = 1.405 + 0.47 \cdot \log K_{ow}$	Triazoles

Sorption of pH-dependent non-hydrophobic PPCPs

Table 1
Chemical properties, structures and speciation of the sorbates.

Compound (Detection, wavelength λ)	Structure	pK_a^a (character ^c)	$\log K_{OW}^a$	Species ^d at pH =		
				4	6	8
Atenolol ($\lambda = 227$ nm)		9.2 ± 0.4^e (B)	0.1 ± 0.28	+	+	+, 0
Carbamazepine ($\lambda = 232$ nm)		-0.49 ± 0.2^e (B)	2.67 ± 0.38	0	0	0
Diazepam ($\lambda = 232$ nm)		3.4 ± 0.1^e (B)	2.96 ± 0.55	+, 0	0	0
Naproxen ($\lambda = 215$ nm)		4.8 ± 0.3 (A)	3.0 ± 0.24	0, -	-	-
Phenobarbital ($\lambda = 243$ nm)		7.6 ± 0.1 (A)	0.53 ± 0.26	0	0	-
Primidone ($\lambda = 215$ nm)		12.3 ± 0.4 (A)	0.4 ± 0.52	0	0	0
Trimethoprim ($\lambda = 225$ nm)		7.2 ± 0.1^e (B)	0.79 ± 0.38	+	+	+, 0











Fig. 1. Chemical structures and conjugated species of phenol, o-phthalic acid, and nicotinic acid used in this study. The pK_a values indicated are those in pure water (pK_a^w).

Fabrice Gritti (2009)

Sorption of pH-dependent non-hydrophobic PPCPs

Table 1
Chemical properties, structures and speciation of the sorbates.

Compound (Detection, wavelength λ)	Structure	pK_a^a (character ^c)	$\log K_{OW}^a$	Species ^d at pH =		
				4	6	8
Atenolol ($\lambda = 227$ nm)		9.2 ± 0.4^e (B)	0.1 ± 0.28	+	+	+, 0
Carbamazepine ($\lambda = 232$ nm)		-0.49 ± 0.2^e (B)	2.67 ± 0.38	0	0	0
Diazepam ($\lambda = 232$ nm)		3.4 ± 0.1^e (B)	2.96 ± 0.55	+, 0	0	0
Naproxen ($\lambda = 215$ nm)		4.8 ± 0.3 (A)	3.0 ± 0.24	0, -	-	-
Phenobarbital ($\lambda = 243$ nm)		7.6 ± 0.1 (A)	0.53 ± 0.26	0	0	-
Primidone ($\lambda = 215$ nm)		12.3 ± 0.4 (A)	0.4 ± 0.52	0	0	0
Trimethoprim ($\lambda = 225$ nm)		7.2 ± 0.1^e (B)	0.79 ± 0.38	+	+	+, 0

Sorption of pH-dependent non-hydrophobic PPCPs at pH = 8.23

Table 3

Sorption model variables and model fitting adjustments.

Compound	Character (charge)	Linear		Freundlich			Langmuir			HI
		K_d	STD	K_F	K_L	C_{max}	R^2	n	R^2	
CBZ	Base (N)	0.40	0.55	0.97	0.34	5.62	0.88	0.37	0.97	–
ACP	Acid (N)	0.50	0.16	0.68	0.02	32.93	0.98	0.87	0.97	–
ATN	Base (+)	7.93	4.01	11.00	0.08	189.01	0.99	0.76	0.99	–0.66
CAF	Base (+)	17.86	6.10	20.00	0.09	221.17	0.98	0.87	0.99	–0.30
NPX	Acid (–)	1.86	1.27	3.38	0.07	68.68	0.97	0.70	0.99	4.46
SX7	Acid (–)	4.25	1.61	6.20	0.03	218.78	1.00	0.79	0.99	13.81


K_d expressed in $L\ kg^{-1}$; K_F expressed in $\mu g^{1-n} L^n\ kg^{-1}$; K_L expressed in $L\ \mu g^{-1}$; C_{max} expressed in $\mu g\ kg^{-1}$. STD: Standard deviation. HI: Hysteresis Index.

N: Neutral (+): Positively charged (–): Negatively charged.

Martinez-Hernandez (2014)

- Basic compounds : Highest K_d by strong interactions with negatively-charged surface
- Acidic compounds : Moderate K_d by ion exchange with cations on surface or physical interaction among polar molecules
- Neutral compounds : Lowest K_d by negligible sorption affinity

K_d in sludges



K_d in sludges

German WWTP consists of a biological treatment unit with two denitrifying tanks, a nitrifying tank and a secondary clarifier with the sludge recycle going to the inlet of the first denitrification tank

	Location	TOC/TSS	COD/TSS	P/TSS	Fe(III)/TSS	N/TSS	VSS/TSS
Primary	After Grit	35%	146%	3%	<1%		
Secondary	Nitrification tank	34%	112%	3%	4%	5.8%	60%

Table 1.3: Sorption coefficients for sludge of municipal wastewater treatment, measured in batch experiments and octanol-water partitioning coefficients (according to Ternes *et al.* 2004).

	Primary sludge K_d [L kg _{SS} ⁻¹]	Secondary sludge K_d [L kg _{SS} ⁻¹]	log K_{OW}
	I.	II.	III.
Acidic pharmaceuticals			
Diclofenac	459±32	16.0±3.1	4.6
Ibuprofen	-- (< 20)	7.1±2.0	3.5
Clofibric acid	-- (< 30)	4.8±2.5	2.57
Neutral pharmaceuticals			
Ifosfamide	21.8±13.8	1.4±0.4	0.86
Cyclophosphamide	55.4±19.6	2.4±0.5	0.63
Carbamazepine	-- (< 20)	1.2±0.5	2.45
Diazepam	43.9±26.1	21.1±7.6	2.82
Musk fragrances			
Galaxolide (HHCB)	4919±2073	1807±534	5.9
Tonalide (AHTN)	5299±1905	2372±958	5.7
Iodinated contrast media			
Iopromide	-- (<5)	11±1	-2.33
Estrogens			
17α-Ethinylestradiol	278±3	349±37	3.9


Removal of PPCPs in WWTP

Biological removal (g day⁻¹)

Level of removal
by biodegradation^a

Day 1 Day 3 Day 9 Day 10 Average Standard deviation

Diclofenac	-23.1	-4.1	-47.5	-24.1	-24.7	17.8	Low
Etofenamate	-0.3 ^b	-1.1	-2.6	12.4	2.1	6.9	Medium
Ibuprofen	45.1	0.5	0.4	-0.3	11.4	22.5	High
Ketoprofen	83.3	-0.2					
Fluoxetine	0.0 ^b	0.0 ^b					
Clorazepate	0.0 ^b	0.2					
Hydroxyzine	-2.3 ^b	-1.6 ^b					
Indapamide	-0.9 ^b	-0.1 ^b					
Enalapril	0.0 ^b	-0.3					
Captopril	2.1	-0.6					
Atenolol	12.8	2.8					
Clofibric acid	49.9	0.2					
Ampicillin	-0.1 ^b	0.0 ^b					
Galaxolide	-0.4	-3.7					
Tonalide	0.2	-0.3					
Cashmeran	-0.8	-3.8					
Celestolide	0.5	-0.3					
Traseolide	0.2	-0.1					



Biological
Adsorption
UV

	Standard deviation	Theoretical logK _{ow}	Level of removal by adsorption ^a
	7.3	4.0–4.7	Medium
	1.3	4.2	Medium
	0.5	3.5–4.0	Low
	0.1	3.0–3.2	Low
	0.0	1.8	Low
	0.4	2.1	Low
	1.3	2.4	Medium
	3.2	2.7	High
	0.2	2.4	Low
	0.1	1.7	Low
	0.0	0.2–1.4	Low
	0.3	2.6–3.6	Low
	0.0	1.4	Low
	7.3	4.6–5.9	High
	1.0	4.8–5.7	High
	1.2	4.8	High
	0.1	4.4	High
	0.0	6.3	High

R. Salgado (2011)

Any questions?

To next lecture

