

[Print version](#)

CEE 697z

Organic Compounds in Water and Wastewater

Cyanotoxins
Compounds, Toxicity and Occurrence


Lecture #27

CEE 697z - Lecture #27


Cyanotoxins

► Neurotoxins


- Anatoxin-a
- Anatoxin-as
- Saxitoxin
- β -Methylamino-L-alanine (BMAA)




Anatoxin-a



Anatoxin-as




Saxitoxin




BMAA

► Hepatotoxins

- Microcystins
- Nodularins



Microcystin LN



Nodularin R

► Cytotoxins


- Cylindrospermopsin



Cylindrospermopsin



► Gastrointestinal and dermatotoxins

- Aplysiatoxin
- Lyngbyatoxin a




Aplysiatoxin

CEE 697z - Lecture #27




Saxitoxins

- ▶ Saxitoxin is usually associated with red tides in marine ecosystems
 - ▶ Responsible for paralytic shellfish poisoning
 - ▶ Been detected in some freshwater species



Saxitoxin



Toxin	R ₁	R ₂	R ₃	R ₄	R ₅
STX	H	H	H	CONH ₂	OH
GTX2	H	H	OSO ₃ ⁻	CONH ₂	OH
GTX3	H	H	H	CONH ₂	OH
GTX5	H	H	H	CONHSO ₃ ⁻	OH
C1	H	H	OSO ₃ ⁻	CONHSO ₃ ⁻	OH
C2	H	H	OSO ₃ ⁻	CONHSO ₃ ⁻	OH
C3	OH	H	OSO ₃ ⁻	CONHSO ₃ ⁻	OH
C4	OH	OSO ₃ ⁻	H	CONHSO ₃ ⁻	OH
neoSTX	OH	H	H	CONH ₂	OH
GTX1	OH	H	OSO ₃ ⁻	CONH ₂	OH
GTX4	OH	H	OSO ₃ ⁻	CONH ₂	OH
GTX6	OH	H	H	CONHSO ₃ ⁻	OH
dcSTX	H	H	H	H	OH
dcneoSTX	OH	H	H	H	OH
dcGTX1	OH	H	OSO ₃ ⁻	H	OH
dcGTX2	H	H	OSO ₃ ⁻	H	OH
dcGTX3	H	OSO ₃ ⁻	H	H	OH
dcGTX4	OH	OSO ₃ ⁻	H	H	OH
LWTX1	H	OSO ₃ ⁻	H	COCH ₃	H
LWTX2	H	OSO ₃ ⁻	H	COCH ₃	H
LWTX3	H	H	OSO ₃ ⁻	COCH ₃	H
LWTX4	H	H	H	H	H
LWTX5	H	H	H	COCH ₃	OH
LWTX6	H	H	H	COCH ₃	H

CEE 697z+ Lecture #27

Saxitoxins




Fig. 4 Saxitoxin analogues produced by some members of different cyanobacteria genera.


Araoz, R., Molgo, J. and de Marsac, N.T. (2010)
Neurotoxic cyanobacterial toxins. *Toxicon* 56(5), 813-828.

CEE 697z - Lecture #27

BMAA neurotoxin

- Caused by over 30 species of cyanobacteria:
 - Ex. *Microcystis, Anabaena, Nostoc, Planktothrix*
- Can cause motor neuron disease or death
- Accumulates in brain tissue
- Found in Guam and linked to ALS

β -methylamino-L-alanine



CEE 697z - Lecture #27

Amyotrophic Lateral Sclerosis (ALS)

- Neurodegenerative disease
- About 2 per 100,000 people in US
- Can be caused by the neurotoxin BMAA
- Symptoms
 - Muscle weakness (including speech muscles)
 - Twitching and cramping of muscles
 - Trouble with speech
 - Shortness of breath, trouble swallowing
 - Death by suffocation

CEE 697z - Lecture #27

Parkinson's Disease (PD)

- Neurodegenerative Disease
- Symptoms
 - Rigidity of muscles, slowing of movement
 - Muscle spasms or tremors
 - Loss of smell, blinking, smiling
 - Speech changes (soft, monotone, repetition)
 - Dementia in later stages

CEE 697z - Lecture #27

Alzheimer's


- 7th leading cause of death
- Most common form of dementia
- Destroys brain cells leading to memory loss, confusion, changes in personality, mood, behavior, problems with language

CEE 697z - Lecture #27

Hepatotoxins


- Cyclic peptides
- Cause liver damage
- Long term exposure can lead to liver cancer

Merel, S., Walker, D., Chicana, R., Snyder, S., Baures, E. and Thomas, O. (2013) State of knowledge and concerns on cyanobacterial blooms and cyanotoxins. Environment International 59, 303-327.



Microsystins

- ▶ Polypeptide produced by **Microsystis & others**
 - ▶ Adda is: 3-amino-9-methoxy-2,6,8-trimethyl-10-phenyldeca-4,6-dienoic acid
- ▶ 90 congeners & 200 related compounds
- ▶ LD₅₀ ~25-60 ug/kg (cyanide is 4,000 ug/kg)
- ▶ Hepatotoxin and tumor promoter



Nodularins

- ▶ Powerful hepatotoxins
- ▶ Cyclic nonribosomal peptide

- ▶ Similar to microcystins, as both have 3-amino-9-methoxy-2,6,8-trimethyl-10-phenyldeca-4,6-dienoic acid (Adda)
- ▶ Difference is Nodularins have 2-(methylamino)-2-dehydrobutyric acid (Mdhb) where Microcystins have dehydroalanine

- ▶ Produced by *Nodularia spumigena*, a cyanobacterium

- ▶ The late summer blooms of *Nodularia spumigena* are among the largest cyanobacterial mass occurrences in the world.

Nodularin

Very similar to microcystins, except that nodularins do not bind covalently to proteins in the body and thus move more easily throughout the body and cells






Fig. 1a Structure of microcystins and nodularins.

Merel, S., Walker, D., Chicana, R., Snyder, S., Baures, E. and Thomas, O. (2013) State of knowledge and concerns on cyanobacterial blooms and cyanotoxins. *Environment International* 59, 303-327.



Cylindrospermopsin


- ▶ Alkaloid
- ▶ Produced by Cylindrospermopsis
- ▶ LD₅₀ 300 µg/kg
- ▶ Hepatotoxin and Neurotoxin
- ▶ Subtropical species recently reported in Michigan



CEE 697z - Lecture #27

Cylindrospermopsin: Variants

- ▶ The molecular structures of cylindrospermopsin (1) and its analogs 7-deoxy-cylindrospermopsin (2) and 7-epicylindrospermopsin (3).



Toxicity of Algal Toxins Relative to Other Toxic Compounds found in Water

▶ Reference Dose = amount that can be ingested orally by a person, above which a toxic effect may occur, on a milligram per kilogram body weight per day basis.

Toxin Reference Doses

- ← Dioxin (0.000001 mg/kg-d)
- ← Microcystin LR (0.000003 mg/kg-d)
- ← Saxitoxin (0.000005 mg/kg-d)
- ← PCBs (0.00002 mg/kg-d)
- ← Cylindrospermopsin (0.00003 mg/kg-d)
- ← Methylmercury (0.0001 mg/kg-d)
- ← Anatoxin-A (0.0005 mg/kg-d)
- ← DDT (0.0005 mg/kg-d)
- ← Selenium (0.005 mg/kg-d)
- ← Botulinum toxin A (0.001 mg/kg-d)
- ← Alachlor (0.01 mg/kg-d)
- ← Cyanide (0.02 mg/kg-d)
- ← Atrazine (0.04 mg/kg-d)
- ← Fluoride (0.06 mg/kg-d)
- ← Chlorine (0.1 mg/kg-d)
- ← Aluminum (1 mg/kg-d)
- ← Ethylene Glycol (2 mg/kg-d)

CEE 697z - Lecture #27

US Regulatory Action

From: Cyanobacteria and Cyanotoxins: Information for Drinking Water Systems , USEPA , Ju

Table 1. Cyanotoxins on the Contaminant Candidate List (CCL)

Cyanotoxin	Number of known variants or analogues	Primary organ affected	Health Effects ¹	Most common Cyanobacteria producing toxin ²
Microcystin-LR	80-90	Liver	Abdominal pain Vomiting and diarrhea Liver inflammation and hemorrhage Acute pneumonia Acute dermatitis Kidney damage Potential tumor growth promotion	<i>Microcystis</i> <i>Anabaena</i> <i>Planktothrix</i> <i>Anabaenopsis</i> <i>Aphanizomenon</i>
Cylindrospermopsin	3	Liver	Tingling, burning, numbness, drowsiness, incoherent speech, salivation, respiratory paralysis leading to death	<i>Cylindrospermopsis</i> <i>Aphanizomenon</i> <i>Anabaena</i> <i>Lyngbya</i> <i>Raphidiopsis</i> <i>Umezakia</i>
Anatoxin-a group ³	2-6	Nervous System		<i>Anabaena</i> <i>Planktothrix</i> <i>Aphanizomenon</i> <i>Cylindrospermopsis</i> <i>Oscillatoria</i>

Source: Harmful Algal Research and Response National Environmental Science Strategy (HARRNESS)
² Not all species of the listed genera produce toxin; in addition, listed genera are not equally as important in producing cyanotoxins.
³The anatoxin-a group does not include the organophosphate toxin anatoxin-aSj as it is a separate group. In the US, the most common member is thought to be anatoxin-a, and thus this toxin is listed specifically.

CEE 697z - Lecture #27





TABLE 1 Toxigenic cyanobacteria from marine, brackish and freshwaters		
Cyanotoxin	Main producing cyanobacteria	Bibliographic source
Microcystins	Most of <i>Microcystis</i> spp and <i>Planktothrix</i> spp, some <i>Anabaena</i> , <i>Nostoc</i> and <i>Synechocystis</i> and <i>Cyanothrix bacillare</i> , <i>Arthrospira fusiformis</i> , <i>Limnothrix redekei</i> , <i>Phormidium formosum</i> , <i>Hapalosiphon hibernicus</i>	Sivonen and Jones, 1999; Cronberg et al., 2003; Odebrecht et al., 2002; Ballot et al., 2004; Gkelis et al., 2001; Steffensen et al., 2001; Prinsen et al., 1992
Nodularins	<i>Nodularia spumigena</i> (in transitional waters)	Rinheart et al., 1988
Cylindrospermopsin	<i>Cylindrospermopsis raciborskii</i> , <i>Umezakia natans</i> , <i>Aphanizomenon ovalisporum</i> , <i>Aphanizomenon flos-aquae</i> , <i>Raphidiopsis curvata</i> , <i>Anabaena lapponica</i> , <i>Anabaena bergeri</i>	Ohtani et al., 1992; Harada et al., 1994; Banker et al., 1997; Schemberg et al., 2001; Li et al., 2001; Fastner et al., 2007; Spoof et al., 2006
Anatoxin-a	Most of <i>Anabaena</i> spp., some <i>Aphanizomenon</i> (<i>A. flos-aquae</i> , <i>A. issatschenkoi</i>), <i>Cylindrospermum</i> , <i>Microcystis</i> and <i>Planktothrix</i> spp. and <i>Raphidiopsis mediterranea</i>	Edwards et al., 1992; Sivonen et al., 1989; Park et al., 1993; Namikoshi et al., 2003; Wood et al., 2007
Homoanatoxin-a	<i>Oscillatoria formosa</i> , <i>Raphidiopsis mediterranea</i>	Skulberg et al., 1992; Steffensen et al., 2001; Namikoshi et al., 2003
Anatoxin a-(s)	<i>Anabaena flos-aquae</i> and <i>A. lemmermannii</i>	Carmichael and Gorham, 1978; Henriksen et al., 1997
Saxitoxins (PSP)	<i>Aphanizomenon</i> , <i>Anabaena</i> , <i>Lyngbya</i> and <i>Cylindrospermopsis</i> spp.	Humpp et al., 1994
LPS endotoxins	All cyanobacteria	McElhiney and Lawton, 2005
Aplysiatoxin, Lyngbyatoxin Debromoaplysiatoxin	<i>Lyngbya majuscula</i> (marine waters), <i>Oscillatoria nigro-viridis</i>	Serdula et al., 1982; Mynderse et al., 1997
Microviridin J	<i>Microcystis</i> spp	Rohrlack et al., 2003
β -N-methylamino-L-alanine	<i>Microcystis</i> , <i>Anabaena</i> , <i>Nostoc</i> and <i>Planktothrix</i> spp and most of cyanobacteria symbionts tested	Cox et al., 2005

Funari E, Testai E. Toxigenic cyanobacteria from marine, brackish and freshwaters. Chart. *Critical Reviews in Toxicology*, Feb 2008; 38(2):98 Available from: Academic Search Premier, Ipswich, MA. Accessed March 20, 2010.

TABLE 3 Main toxicological data of some cyanotoxins						
Cyanotoxin	i.p. LD ₅₀ ($\mu\text{g}/\text{kg}$ b.w.)	Oral LD ₅₀ ($\mu\text{g}/\text{kg}$ b.w.)	Target organ and mechanism of action	NOEL ($\mu\text{g}/\text{kg}/\text{d}$)	LOEL ($\mu\text{g}/\text{kg}/\text{d}$)	ADI/TDI ($\mu\text{g}/\text{kg}/\text{d}$)
Microcystin (MC)	50–1200	5000	Liver (PP1 and PP2A phosphatases inhibition-Tumor promoting activity)	40 (MC-LR; mice; 13 weeks; gavage) 350 (MC-LR in BGAS extracts; mice; 13 weeks; dietary) ND (refer to MC-LR)	200 ND	0.04 — (UF = 1000)
Nodularin (NOD)	50	ND	Liver (PP1 and PP2A phosphatases inhibition-Tumor promoting activity)	ND	—	—
Cylindrospermopsin (CYN)	2100 (24 h) 200 (6 days)	4400–6900 (2–6 days)	Kidney, liver (Parent compound: protein synthesis inhibition; Metabolites: different but unknown mechanism; possible genotoxicity)	30 (Mice; 11 weeks; gavage) <i>C. raciborskii</i> extracts more toxic than pure CYN	60	0.03 (UF = 1000)
Anatoxin-a	375	5000	Neuromuscular system (Depolarizing effect due to binding to nicotinic Ach receptor)	>510 (mice; 54 days; drinking water) Limited chronic risk	ND	0.51 (UF = 1000)
Homoanatoxin-a	330	ND	Similar to anatoxin-a	ND Limited chronic risk	—	—
Anatoxin a-(s)	20–40		Peripheral nervous system (AChE inhibition; nerve hyper-excitability)	ND Limited chronic risk	—	—
Saxitoxin (STX)	10–20	263	Neuromuscular system (Membrane ion channel block) Human: 0.144–0.304 mg/person: mild symptoms 0.456–12 mg/person: from moderate symptoms up to paralysis and death	ND Acute risk > chronic	—	—
LPS Endotoxins	40–190 mg/kg bw	ND	Skin and mucosa (irritation, topical effects)	ND	—	—

i.p. = intraperitoneal; ND = Not determined; UF = uncertainty factor.
Note: bibliographic references are available within the text.

Funari E, Testai E. Toxigenic cyanobacteria from marine, brackish and freshwaters. Chart. *Critical Reviews in Toxicology*, Feb2008; 38(2): 101 Available from: Academic Search Premier, Ipswich, MA. Accessed March 20, 2010. CEE 697z - Lecture #27




Lake Erie; western basin



The 2014 Toledo Ohio incident:

- ▶ On-line reports
- ▶ http://www.nytimes.com/2014/08/05/us/lifting-ban-toledo-says-its-water-is-safe-to-drink-again.html?_r=0
- ▶ <http://www.vox.com/2014/8/3/5963645/a-toxic-algae-bloom-has-left-400000-people-in-ohio-without-drinking>


CEE 497s - Lecture #27



Microcystin Concentrations


- ▶ **1 ppb WHO drinking water limit**
- ▶ **20 ppb WHO swimming limit**
- ▶ **60 ppb highest level for Lake Erie till 2011**
- ▶ **84 ppb highest level for Grand Lake St. Marys till 2010**
- ▶ **2000+ Grand Lake St. Marys 2010**
- ▶ **1200 Lake Erie Maumee Bay area 2011**
- ▶ **Carroll Water System, west of Davis-Besse, 4&5 Sept 2013, 1.4 and 3.5 ppb**

CEE 697z - Lecture #27



Sampling locations and microcystin-LR concentrations (average \pm S.E., $\mu\text{g/L}$) along the Lake Ontario shoreline and the associated rivers, embayments and ponds. The green vertical bar at each Lake Ontario site represents the "shoreside" sampling site. The red vertical bar at each Lake Ontario site represents samples taken in the creek, river or embayment. Vertical bars are to scale. Bars for Knauf and Georgetown Ponds, Lake Neatahwanta, Sackets Harbor "scum" and the Conesus and Silver Lake "scum" concentrations are not to scale with concentration listed above the bar. The vertical bars for the nearshore and offshore of Lake Ontario are labeled "30 m" and "100 m".

CEE 697z - Lecture #27



Average monthly ($\pm\text{S.E.}$) microcystin-LR, total phosphorus and phycocyanin concentration ($\mu\text{g/L}$) at 37 sites in Lake Ontario from 2003–2006. Sites include streams, rivers, embayments, shoreside sites, and the nearshore and offshore zones. See [Fig. 1](#) for location of sites.

CEE 697z - Lecture #27

► [To next lecture](#)

CEE 697z - Lecture #27