

IV. CALCULUS OF VECTORS, DYADICS & TENSORS

A. Introduction & Review

1. In this section we will review some elementary ideas of vector calculus which you have seen in earlier math, physics and mechanics (statics or dynamics) courses. We will then try to develop a more 'user-friendly' approach to the study of vector calculus and introduce the concept of tensors.

In turn we will discuss

- scalar (or dot or inner) product
- vector (or cross) product
- index notation and the summation convention for making vector calculations fast & easy
- vector calculus \rightarrow taking derivatives of scalar and vector functions
 \rightarrow gradient, divergence, curl
- Lots of vector operations and identities
- Integral Theorems - Divergence and Stokes Theorem.
 Also, different forms of Green's Theorem, which will arise again in our study of pdes.
- Application of some of these ideas to derive a diffusion eqn, study geometry of curves (and surfaces), etc.
- vector calculus of orthogonal curvilinear coordinates such as cartesian, cylindrical and spherical coordinates, as well as a discussion of the general case which leads to such systems as oblate and prolate spheroidal coordinates.
- an introduction to tensors and dyadics
 ... and the answer to the question "What is a tensor?"

NOTE: There are a number of books which focus on teaching about vectors, tensors and vector calculus. Several of my favorite references for reading about ideas and examples are
 F. Chorlton, Vector & Tensor Methods
 H.M. Schey, Div, Grad, Curl and all That
 J.G. Simmonds, A Brief on Tensor Analysis

2. Scalars & vectors

a. A scalar is a quantity or function with magnitude only
e.g. mass and temperature are common scalars.

b. A vector is characterized by magnitude and direction.

(i) The most common examples of vectors are a force applied to move a given object (clearly you must know both the magnitude of the force as well as its direction in order to describe what happens to the object) and the velocity with which a given object moves (e.g., if you compare driving in a straightline at 20 mph or driving at 20 mph on a circular track, then you realize in order to completely characterize the motion it is necessary to specify both the magnitude of the velocity - i.e., speed - as well as direction).

(ii) Vectors are typically represented geometrically with arrows  (length denotes magnitude; orientation indicates direction)

(iii) Two vectors are equal if they have the same magnitude and same direction.



conclude $a = b$

sometimes this is called
"parallel transport of vectors"

→ Nevertheless, it is very important to keep in mind that the effect of a given vector may, indeed often will, depend on its location.


(iv) NOTATION: I will typically indicate a vector quantity by an underline: a or b (books use bold-faced type)
(Another common notation is to use arrows: \vec{a} or \vec{b})

one final comment on notation: a UNIT VECTOR by definition has length = 1


2 Cartesian Coordinate System

a. We will describe a vector by giving its components relative to a cartesian, or x, y, z , coordinate system.

b. We will indicate the base vectors as (by definition the base vectors have length = 1)



or



"What you have probably seen"

"What we will use"

NOTE: We will indicate the 3 coordinate directions by x_1, x_2, x_3 where $x_1 \equiv x, x_2 \equiv y, x_3 \equiv z$.

c. In order to describe a vector you must give both the components and the base vectors:

For example: $\underline{q} = q_x \underline{i} + q_y \underline{j} + q_z \underline{k}$ (j14)

or we will write: $\underline{q} = q_1 \underline{e}_1 + q_2 \underline{e}_2 + q_3 \underline{e}_3$
 $\rightarrow \text{length: } |\underline{q}| = \sqrt{q_x^2 + q_y^2 + q_z^2}$

4. SCALAR product (also called the DOT or INNER product)

a. The scalar product of two vectors \underline{q} and \underline{b} is defined as

$$(j15) \quad \underline{q} \cdot \underline{b} = |\underline{q}| |\underline{b}| \cos\theta$$


where $|\underline{q}|$ and $|\underline{b}|$ are the lengths, or magnitudes, of the two vectors



(scalar product continued)

b. The scalar product is commutative and distributive.

$$\underline{a} \cdot \underline{b} = \underline{b} \cdot \underline{a} \quad \text{and} \quad (\underline{a} + \underline{b}) \cdot \underline{c} = \underline{a} \cdot \underline{c} + \underline{b} \cdot \underline{c}$$

c. Two vectors are orthogonal if $\underline{a} \cdot \underline{b} = 0$ 

If \underline{a} and \underline{b} are both nonzero vectors ($\underline{a}, \underline{b} \neq \underline{0}$)
then $\underline{a} \cdot \underline{b} = 0 \rightarrow \theta = \pi/2$ and the two vectors are perpendicular.

d. Since the cartesian unit vectors $\underline{i}, \underline{j}, \underline{k}$ satisfy

$$\underline{i} \cdot \underline{i} = 1, \underline{i} \cdot \underline{j} = 0, \underline{i} \cdot \underline{k} = 0, \text{ etc. then}$$

$$\underline{a} \cdot \underline{b} = a_x b_x + a_y b_y + a_z b_z \quad (116)$$

$$= a_1 b_1 + a_2 b_2 + a_3 b_3 \quad \text{in our new notation.}$$


$$\text{Also, } \underline{a} \cdot \underline{a} = a_1^2 + a_2^2 + a_3^2 = |\underline{a}|^2 \quad (117) \quad (\text{square of the length})$$

5. VECTOR (or cross) product of two vectors

d. The vector product of 2 vectors \underline{a} and \underline{b} is defined as

NOTE
My notation
for this operation
is $\underline{a} \wedge \underline{b}$; many others
write \times

$$\underline{a} \wedge \underline{b} = |\underline{a}| |\underline{b}| \sin \theta \underline{e} \quad (118)$$



where \underline{e} is a unit vector in the direction perpendicular to the plane formed by \underline{a} and \underline{b} as given by the RIGHT-HAND RULE.

b. From the definition: $\underline{a} \wedge \underline{a} = \underline{0}$ (cross product of a vector with itself = $\underline{0}$)Important
to understand
this

$$\text{Also, } \underline{a} \wedge \underline{b} = - \underline{b} \wedge \underline{a} \quad (119a) \quad \text{and} \quad \underline{a} \wedge (\underline{b} \wedge \underline{c}) = \underline{a} \wedge \underline{b} + \underline{a} \wedge \underline{c} \quad (119b)$$

If follows that $\underline{i} \wedge \underline{j} = \underline{k}$; $\underline{j} \wedge \underline{k} = \underline{i}$; $\underline{k} \wedge \underline{i} = \underline{j}$

c. You may also remember writing something like

$$\underline{a} \wedge \underline{b} = \det \begin{vmatrix} \underline{i} & \underline{j} & \underline{k} \\ a_x & a_y & a_z \\ b_x & b_y & b_z \end{vmatrix} = \underline{i} (a_y b_z - a_z b_y) + \underline{j} (a_z b_x - a_x b_z) + \underline{k} (a_x b_y - a_y b_x) \quad (119c)$$

+ rather laborious
to write!

Much of the above is cumbersome and frightfully lengthy to write. We now introduce a convenient shorthand notation which will simplify many manipulations.

B. EINSTEIN INDEX NOTATION & THE SUMMATION CONVENTION

1. Let us reconsider^{some of} the above manipulations. From now on keep in mind that we are typically having to represent vectors in a three-dimensional world and so we will always write vectors with 3 components.

2. Label the (x_1, y_1, z_1) coordinates by $(1, 2, 3)$.

a. Denote the vector \underline{q} with components q_i and base vectors \underline{e}_i , where the subscript i may take on the 3 values $i=1, 2$ or 3 . We write

$$(120) \quad \underline{q} = q_1 \underline{e}_1 + q_2 \underline{e}_2 + q_3 \underline{e}_3 = \sum_{i=1}^3 q_i \underline{e}_i \equiv q_i \underline{e}_i \quad (= q_j \underline{e}_j)$$

^{↑ dummy summation symbol}

IMPORTANT:

b. From now on we will not usually write the summation symbol. Instead we will invoke the **SUMMATION CONVENTION** - if an index (i) appears **TWICE** in an expression, then we will know that we should perform a summation with $i=1, 2$ and 3 .

SUMMATION CONVENTION

Thus, the vector \underline{q} is expressed as $\underline{q} = q_i \underline{e}_i = q_m \underline{e}_m$


→ This idea of implied summation when an index appears **TWICE** must be clear in your mind before you move on.

[↑]
dummy summation
index; two sides
of equal sign mean
the same thing.

c. This notation simply allows us to proceed without carrying around the summation symbol.

) 3. inner product of base vectors and the Kronecker delta symbol 128

a. We previously saw that $\underline{i} \cdot \underline{i} = 1$, $\underline{j} \cdot \underline{j} = 1$, $\underline{i} \cdot \underline{j} = 0$, $\underline{i} \cdot \underline{k} = 0$, etc or in terms of the $(1, 2, 3)$ notation we may write



$$\underline{e}_1 \cdot \underline{e}_1 = 1 ; \underline{e}_2 \cdot \underline{e}_2 = 1 ; \underline{e}_3 \cdot \underline{e}_1 = 0 , \underline{e}_1 \cdot \underline{e}_3 = 0 , \underline{e}_2 \cdot \underline{e}_3 = 0 , \text{etc}$$

b. Thus, we see that

$$(121) \quad \underline{e}_i \cdot \underline{e}_j = \begin{cases} 1 & \text{if } i=j \\ 0 & \text{if } i \neq j \end{cases} \quad \begin{matrix} \text{orthogonality} \\ \text{of the} \\ \text{base vectors} \end{matrix}$$

where i and j may take on any of the values 1, 2 or 3.

↳ a convenient shorthand notation

c. Kronecker delta δ_{ij}

let us define the symbol δ_{ij} as

$$(122) \quad \delta_{ij} = \begin{cases} 0 & i \neq j \\ 1 & i=j \end{cases} \quad \begin{matrix} \text{where } i=1,2,3 \\ j=1,2,3 \end{matrix}$$

Therefore we see that

$$(123) \quad \underline{e}_i \cdot \underline{e}_j = \delta_{ij} \quad \begin{matrix} (\text{which } = 1 \text{ if } i=j) \\ \text{and } = 0 \text{ if } i \neq j \end{matrix}$$

4. The scalar product revisited

$$a. \quad \underline{a} \cdot \underline{b} = \sum_{i=1}^3 a_i \underline{e}_i \cdot \sum_{j=1}^3 b_j \underline{e}_j = \sum_{i=1}^3 a_i b_i = a_1 b_1 + a_2 b_2 + a_3 b_3$$

$\uparrow \quad \uparrow$
 $\underline{e}_i \cdot \underline{e}_j = 1 \text{ if } i=j$
 $= 0 \text{ if } i \neq j$

where we have
 invoked the
 summation convention

NOTE: in the second sum, $\sum_{j=1}^3 b_j \underline{e}_j$ we have used the dummy summation index j so as not to confuse operations with the first sum utilizing i as a summation index

b. Use the δ_{ij} shorthand and the summation convention, *only numbers if $i=j$*

$$\underline{a} \cdot \underline{b} = a_i \underline{e}_i \cdot b_j \underline{e}_j = a_i b_j (\underline{e}_i \cdot \underline{e}_j) = a_i b_j \delta_{ij} = a_i b_i$$

*vector operations;
only acts on vectors and
not the components.*

*actually represents the
double sum*

$$\sum_{i=1}^3 \sum_{j=1}^3 a_i b_j \delta_{ij}$$

c. $\underline{a} \cdot \underline{b} = a_i b_i (= a_1 b_1 + a_2 b_2 + a_3 b_3)$

(124)

δ_{ij} is sometimes
called the
"replacement operator".

c. **VERY IMPORTANT:** We are using the summation convention to recognize that whenever we see a repeated index (e.g., $a_i \underline{e}_i$, or $a_i b_i$) we mean an implied sum with the index taking the values 1, 2 and 3. This, it is important to use different summation labels when beginning operations: For example,

$$\underline{a} \cdot \underline{b} = a_i \underline{e}_i \cdot b_j \underline{e}_j$$

↑ a different summation index!

*NEVER WRITE $a_i \underline{e}_i \cdot b_i \underline{e}_i$
which is very confusing.*

d. A few remarks

(i) $\delta_{ii} = \delta_{11} + \delta_{22} + \delta_{33} = 3$

where we have used the summation convention.

(ii) δ_{ij} : the replacement operator. $\Rightarrow \delta_{ij} c_j = c_i$

(iii) Sometimes people are sloppy and do not write the unit vectors i.e., instead of $\underline{a} = a_i \underline{e}_i$, occasionally you see written $\underline{a} = a_i$ which you can recognize since an index will appear only once ('free index') whenever performing vector operations I will **ALWAYS** keep track of the unit vectors.

(iv) Example:

$$(\underline{a} \cdot \underline{b}) \underline{c} = a_i b_i c_i = a_i b_i c_j \underline{e}_j \quad \text{and you might call } (a_i b_i) c_j \text{ the } j^{\text{th}} \text{ component of the vector.}$$

*can only take
on a value 1, 2
or 3*

130.

* * this symbol will be useful * *
 whenever vector products arise

5. Permutation Symbol

$$\epsilon_{ijk} \quad i=1,2,3 \quad j=1,2,3 \quad k=1,2,3$$

2. Definition : $\epsilon_{ijk} = \begin{cases} +1 \text{ or } -1 & \text{if } i,j,k \text{ are all different} \\ 0 & \text{if any two indices are the same} \end{cases}$ (125a)

In particular,

(125b) $\epsilon_{ijk} = +1$ if i,j,k are an EVEN permutation of $1,2,3$
 (or $1,2,3$ are in cyclic order)

$$\epsilon_{123} = 1 \quad \epsilon_{312} = 1 \quad \epsilon_{231} = 1$$

(125c) $\epsilon_{ijk} = -1$ if i,j,k are an ODD permutation of $1,2,3$

$$\epsilon_{213} = -1 \quad \epsilon_{132} = -1 \quad \epsilon_{321} = -1$$

NOTE: By even permutation we mean that an even # of interchanges of the indices must occur to get back to the order $1,2,3$; analogous for meaning of odd permutation...

b. This definition has the following cyclic and interchange property:

Compare with
above example:




$$\epsilon_{ijk} = \epsilon_{kij} = \epsilon_{jki} \quad (126a)$$

and if two indices are simply interchanged, the sign changes,

$$(126b) \quad \epsilon_{ijk} = -\epsilon_{ikj} \quad \text{or} \quad \epsilon_{ijk} = -\epsilon_{jik}$$

IMPORTANT

Also, since i,j,k can each independently take on the values $1,2,3$, then ϵ_{ijk} represents 27 quantities.



c. We also have

$$\underline{e}_i \wedge \underline{e}_j = \epsilon_{ijk} \underline{e}_k \quad (127)$$

and by referring to the figure at right,
 everything is ok. i.e. $\underline{e}_1 \wedge \underline{e}_2 = +\underline{e}_3 = \underbrace{\epsilon_{123}}_{+1} \underline{e}_3$, etc.

d. We now have an effective shorthand notation for representing the vector product.

let $\underline{c} = \underline{a} \wedge \underline{b}$; write $\underline{a} = a_i \underline{e}_i$, $\underline{b} = b_j \underline{e}_j$

$$\rightarrow \underline{c} = a_i \underline{e}_i \wedge b_j \underline{e}_j = a_i b_j (\underline{e}_i \wedge \underline{e}_j)$$

(128)

$$\underline{a} \wedge \underline{b} = a_i b_j \epsilon_{ijk} \underline{e}_k$$

← NOTE CAREFULLY THE ORDER OF THE INDICES

or with $\underline{c} = c_k \underline{e}_k$, we have

$$c_k = a_i b_j \epsilon_{ijk} \quad \text{which represents 3 terms for } k=1,2,3$$

EXERCISE: Verify that this is in agreement with the 'matrix' definition.

c. triple scalar product: $\underline{a} \cdot (\underline{b} \times \underline{c})$

Again we are careful to use different dummy indices for each vector so

$$\underline{a} \cdot (\underline{b} \times \underline{c}) = a_i \epsilon_i \cdot (b_j \epsilon_j \wedge c_k \epsilon_k) = a_i \epsilon_i \cdot (b_j c_k \epsilon_{jkl} \epsilon_l)$$

$$= a_i b_j c_k \epsilon_{jkl} \epsilon_i \cdot \overbrace{\epsilon_l}^{\delta_{il}}$$

$$= \epsilon_{jki} a_i b_j c_k = \epsilon_{ijk} a_i b_j c_k = (\underline{a} \cdot \underline{b}) \cdot \underline{c} = (\underline{c} \cdot \underline{a}) \cdot \underline{b}$$

Recall also that

$$\underline{a} \cdot (\underline{b} \times \underline{c}) = \det \begin{vmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix} = \epsilon_{ijk} a_i b_j c_k$$

Index representation of the 3×3 determinant.

by using cyclic property of ϵ_{ijk} . Exercise -

convince yourself that these last 2 identities follow from index expression.

5. Useful identities involving ϵ and δ

You are not expected to memorize this eqn. It will be frequently used however.

$$(129) \quad \epsilon_{ijk} \epsilon_{klm} = \delta_{il} \delta_{jm} - \delta_{im} \delta_{jl}$$

i, j, l, m can each independently take on values 1, 2, 3. Hence, this can corresponds to 81 quantities.

Proof: verify by brute force for each of the 81 eqns!

However, it is best to make your life easier by noticing that both sides change sign if either $i \leftrightarrow j$ or $l \leftrightarrow m$ are interchanged. Also, both sides vanishes if $i=j$ or $l=m$. Then, consider remaining terms like:

$$\epsilon_{12k} \epsilon_{k12} = \underbrace{\epsilon_{12j} \epsilon_{j12}}_0 + \epsilon_{132} \epsilon_{212} + \epsilon_{123} \epsilon_{312} = 1$$

and $\delta_{11} \delta_{22} - \delta_{12} \delta_{21} = 1$ so o.k.

Likewise

$$\epsilon_{12k} \epsilon_{k13} = \epsilon_{12i} \epsilon_{i13} + \epsilon_{122} \epsilon_{213} + \epsilon_{123} \epsilon_{313} = 0 \text{; also } \delta_{11} \delta_{23} - \delta_{13} \delta_{21} = 0 \text{ so o.k.}$$

Example 1: Show that $\epsilon_i = \frac{1}{2} \epsilon_{mnj} \epsilon_m \wedge \epsilon_n \delta_{nj} - \delta_{nj} \delta_{ni}$

$$\text{Well, } \epsilon_{mnj} \epsilon_m \wedge \epsilon_n = \epsilon_{mj} \epsilon_{mnj} \epsilon_j = \overbrace{\epsilon_{mj} \epsilon_{mnj}}^0 \epsilon_j = (3 \delta_{ij} - \delta_{jj}) \epsilon_j = 2 \epsilon_i \quad \times$$

Example 2: Show that $\underline{a} \wedge (\underline{b} \times \underline{c}) = \underline{b}(\underline{a} \cdot \underline{c}) - \underline{c}(\underline{a} \cdot \underline{b})$

BAC-CAB identity

$$\begin{aligned} \underline{a} \wedge (\underline{b} \times \underline{c}) &= a_i \epsilon_i \wedge (b_j \epsilon_j \wedge c_k \epsilon_k) = a_i \epsilon_i \wedge (b_j c_k \epsilon_{jkl} \epsilon_l) = a_i b_j c_k \epsilon_{jkl} (\epsilon_i \wedge \epsilon_l) \\ &= a_i b_j c_k \epsilon_{jkl} \epsilon_{ilm} \epsilon_m = a_i b_j c_k \epsilon_{jml} \epsilon_{ilm} \epsilon_m = a_i b_j c_k (\delta_{jm} \delta_{li} - \delta_{jl} \delta_{mi}) \epsilon_m \\ &= a_i b_m c_i \epsilon_m - a_i b_i c_m \epsilon_m = (a_i c_i) b_m \epsilon_m - (a_i b_i) c_m \epsilon_m = (\underline{a} \cdot \underline{c}) \underline{b} - (\underline{a} \cdot \underline{b}) \underline{c} \quad \times \end{aligned}$$

additional

Some Examples of the use of index notation

First, a brief summary of the important ideas

$$(i) \quad e_i \cdot e_j = \delta_{ij} = \begin{cases} 0 & i=j \\ 1 & i \neq j \end{cases}$$

$$(ii) \quad e_i \cdot e_j \cdot e_k = \epsilon_{ijk} e_k \quad \epsilon_{ijk} = \begin{cases} +1 & i,j,k \text{ an even permutation of } 1,2,3 \\ -1 & i,j,k \text{ an odd permutation of } 1,2,3 \\ 0 & \text{any two indices the same} \end{cases}$$

(iii) Summation convention : whenever a subscript appears twice,
a summation from $i=1$ to 3 is implied.
(a subscript should never appear more than twice.)

Examples :

$$(i) \quad \delta_{ik} \delta_{jk} = \delta_{ij} \quad \text{since } \delta_{jk} \text{ is only nonzero when } j=k \text{ so the } k \text{ in } \delta_{ik} \text{ may be replaced by } j.$$

$$(ii) \quad \delta_{ii} = \delta_{11} + \delta_{22} + \delta_{33} = 1 + 1 + 1 = 3 \quad \text{note: Since } i \text{ was a dummy index, } \delta_{ii} = \delta_{kk} = \delta_{mm} \text{ etc.}$$

$$(iii) \quad \delta_{ij} \epsilon_{ijk} = \epsilon_{ijk} = 0 \quad \text{since two of the indices are the same.}$$

$$(iv) \quad \underbrace{\epsilon_{ijk} \epsilon_{nlk}}_{\text{Next, we use the identity}} = \epsilon_{ijk} \epsilon_{klm} \quad \text{by first rotating the indices on the second } \epsilon.$$

$$\epsilon_{ijk} \epsilon_{klm} = \delta_{il} \delta_{jm} - \delta_{in} \delta_{jl}$$

$$\Rightarrow \epsilon_{ijk} \epsilon_{nlj} = \underbrace{\delta_{in} \delta_{jj}}_3 - \delta_{ij} \delta_{nj} = 3 \delta_{in} - \delta_{in} = \underline{2 \delta_{in}}$$

$$(v) \quad a_m b_n \epsilon_{mnq} - a_n b_m \epsilon_{mnq} = ?$$

$\rightarrow m, n$ appear twice in each term so summation is implied.

But, $m \neq n$ are simply dummy variables, i.e., we could just as well use another letter.

So, examine the second term $a_n b_m \epsilon_{mnq}$.

$$a_n b_m \epsilon_{mnq} = -a_n b_m \epsilon_{nmq}; \text{ now let } j=n, m=k$$

$$= -a_j b_k \epsilon_{jknq} \quad \text{which is the same as the above since summation on } j, k \text{ is implied.}$$

$$= -a_m b_n \epsilon_{mnq} \quad \text{by letting } j=m, k=n$$

So, we see that

$$a_m b_n \epsilon_{mnq} - a_n b_m \epsilon_{mnq} = 2 a_m b_n \epsilon_{mnq}$$

$\overbrace{(a_m b_n)}_g \quad g^{\text{th}} \text{ component of } a_m b_n$


→ + DIV, GRAD, CURL.

133.

C. SOME VECTOR CALCULUS (taking derivatives of vector functions)

1. We now wish to consider the derivatives of
 - (i) vector functions which depend on a scalar variable, e.g., calculate rate-of-change of the velocity $\underline{v}(t)$ with respect to time,
 - (ii) scalar functions which depend on position, e.g., the temperature $T(x, y, z)$ at spatial location (x, y, z) ,
 - (iii) vector functions which depend on position, e.g., the electric field $\underline{E}(x, y, z)$ or velocity field of a fluid $\underline{v}(x, y, z)$ at position (x, y, z) .

2. NOTATION: we will typically use the vector \underline{x} to denote the position vector locating a point in space.



One can discuss SCALAR fields $\phi(\underline{x}) = \phi(x_1, x_2, x_3)$ or just ϕ
one can discuss VECTOR fields $\underline{a}(\underline{x}) = a_1(x_1, x_2, x_3) \underline{\epsilon}_1 + a_2(x_1, x_2, x_3) \underline{\epsilon}_2 + a_3(x_1, x_2, x_3) \underline{\epsilon}_3$
each of the components of the vector \underline{a} depends on location in space. \Rightarrow sometimes one sees this written $a_j(t)$

and one can discuss vector fields of a single variable, e.g., $\underline{b}(t)$.

3. Differentiation of vectors

a. Suppose $\underline{q} = \underline{q}(t) = q_i(t) \underline{e}_i$

Cartesian base vectors;
constant length (=1) and
constant direction.

Then, $\frac{d}{dt} \underline{q}(t) = \frac{da_i(t)}{dt} \underline{e}_i$ since the cartesian base
vectors are constant
vectors, $\frac{d}{dt} \underline{e}_i = 0$.

b. By the product rule

$$\frac{d}{dt} [\underline{q}(t) \cdot \underline{b}(t)] = \frac{da}{dt} \cdot \underline{b} + \underline{q} \cdot \frac{db}{dt}$$

c. Exercise: Evaluate $\frac{d}{dt} [\phi(t) \underline{q}(t)]$.

We will now consider spatial derivatives of scalar functions, e.g.,

$$\frac{\partial}{\partial x} \phi(x, y), \frac{\partial \phi}{\partial y}(x, y) \text{ or } \frac{\partial}{\partial x} \underline{b}(x, y, z).$$

4. Gradient operator

Kreysig § 8.8-9; Greenberg § 9.3; Hildebrand § 6.

a. Let $\phi(\underline{x})$ be a scalar function which varies with position x, y, z in space.

The rate of variation of ϕ in the x -direction is $\frac{\partial \phi}{\partial x} = \frac{\partial \phi}{\partial x_1}$, in the
 y -direction is $\frac{\partial \phi}{\partial y} = \frac{\partial \phi}{\partial x_2}$, and in the z -direction is $\frac{\partial \phi}{\partial z} = \frac{\partial \phi}{\partial x_3}$.

b. We introduce the vector

(130)

$$\text{grad } \phi = \nabla \phi = \underline{e}_1 \frac{\partial \phi}{\partial x_1} + \underline{e}_2 \frac{\partial \phi}{\partial x_2} + \underline{e}_3 \frac{\partial \phi}{\partial x_3} = \underline{e}_i \frac{\partial \phi}{\partial x_i}$$

(called the gradient of ϕ)

↑ component in \underline{e}_i is the derivative
with respect to x_i


"commas"
notation to indicate
differentiation with
respect to x_i

$$\text{gradient operator} \equiv \nabla = \underline{e}_1 \frac{\partial}{\partial x_1} + \underline{e}_2 \frac{\partial}{\partial x_2} + \underline{e}_3 \frac{\partial}{\partial x_3} \quad \text{or}$$

$$\nabla = \underline{e}_i \frac{\partial}{\partial x_i} \quad (131)$$

$$\frac{d}{dt} \underline{q}(t) = \lim_{\Delta t \rightarrow 0} \frac{\underline{q}(t + \Delta t) - \underline{q}(t)}{\Delta t}$$

schematic




} schematic interpretation
of the derivative

For those who are interested in a more geometric interpretation of the gradient operator:

Consider a scalar function of position $\phi(\underline{x}) = \phi(x_1, x_2, x_3)$.

Imagine drawing two surfaces in space, labelled S_1 and S_2 , where $\phi = c_1 = \text{constant}$ along S_1 and $\phi = c_2 = \text{constant}$ along S_2



"An exaggerated view of two surfaces along which $\phi(\underline{x}) = \text{constant}$ "

Consider the two nearby points labelled M and N. Clearly, you should recognize and appreciate that since the 'spacing' or distance between the two surfaces changes at different locations along either of the surfaces, then the rate-of-change of ϕ with respect to changes in spatial location will depend on the direction in which the change is made.

Now, for small (infinitesimal) changes, construct a Taylor series about the point M:

$$\phi(N) = \phi(M) + dx_1 \frac{\partial \phi}{\partial x_1} \Big|_M + dx_2 \frac{\partial \phi}{\partial x_2} \Big|_M + dx_3 \frac{\partial \phi}{\partial x_3} \Big|_M + \dots$$

or

$$d\phi = \phi(N) - \phi(M) = d\underline{x} \cdot \nabla \phi$$

Change in ϕ given displacement $d\underline{x}$.

where $\nabla \phi = \sum_i \frac{\partial \phi}{\partial x_i}$
and $d\underline{x} = \sum_j d\underline{x}_j$

5. The directional derivative and the gradient operator

a. Suppose you are interested in some scalar function of position $\phi(\underline{x})$ which may describe the temperature distribution in a component in your stereo or computer. The change in temperature if you are displaced by a small amount $d\underline{x}$ (has magnitude and direction) is given by the (directional derivative):

b. Preliminary: Consider a path, or curve, in space. Measure distance along the curve by s , which is the arclength.

$$\text{where } d\underline{x} = e_i dx_i = e_1 dx_1 + e_2 dx_2 + e_3 dx_3$$

$$(d\underline{x} \cdot d\underline{x}) = |d\underline{x}|^2 = dx_1^2 + dx_2^2 + dx_3^2 = ds$$

Consider a small displacement $d\underline{x}$
where $|d\underline{x}| = ds$ (distance along path)

\Rightarrow The unit tangent vector \underline{t} in the direction of $d\underline{x}$ is

$$\underline{t} = \frac{d\underline{x}(s)}{ds} = e_i \frac{dx_i(s)}{ds}$$

c. Ask how ϕ varies along the curve? $\frac{d\phi}{ds} = ?$

Use the chain-rule:

$$\frac{d}{ds} \phi(x(s)) = \frac{dx_1}{ds} \frac{\partial \phi}{\partial x_1} + \frac{dx_2}{ds} \frac{\partial \phi}{\partial x_2} + \frac{dx_3}{ds} \frac{\partial \phi}{\partial x_3} = \frac{d\underline{x}}{ds} \cdot \nabla \phi$$


position along path

$$(132) \quad \frac{d\phi}{ds} = \underline{t} \cdot \nabla \phi$$

← directional derivative of ϕ in the \underline{t} -direction.

d. Now consider a surface along which $\phi(\underline{x}) = \text{constant}$.

Excuse: Show using index notation that $\underline{t} \cdot \nabla \phi = t_i \frac{\partial \phi}{\partial x_i}$



Now, as $\phi = \text{constant}$ on this surface, then $d\phi = 0$ for any displacement along the surface.

Interpretation of $\nabla \phi$

Then, from (132) $\underline{t} \cdot \nabla \phi = 0$ for movements along the surface. It necessarily follows that $\underline{t} \perp \nabla \phi$...

6. Divergence of a vector field = $\nabla \cdot \underline{f}$ or $\operatorname{div} \underline{f}$

a. Simply compute using standard vceo.

$$\nabla \cdot \underline{f} = (\underline{e}_i \frac{\partial}{\partial x_i}) \cdot (f_j \underline{e}_j) = \underline{e}_i \cdot \frac{\partial f_j}{\partial x_i} \underline{e}_j - f_j \underline{e}_i \cdot \frac{\partial \underline{e}_j}{\partial x_i} \text{ using product rule}$$

$$= \delta_{ij} \frac{\partial f_j}{\partial x_i}$$

common notation sometimes used

$$(133) \quad \nabla \cdot \underline{f} = \frac{\partial f_j}{\partial x_i} = \frac{\partial f_1}{\partial x_1} + \frac{\partial f_2}{\partial x_2} + \frac{\partial f_3}{\partial x_3} = f_{j,j}$$

\downarrow since the \underline{e}_j are unit vectors which do not vary with position in space

b. NOTE: Now that you have gone through this, make your life easier.

The \underline{e}_i are constant vectors with respect to differentiation so we know it is o.k. to simply write

$$\nabla \cdot \underline{f} = \underline{e}_i \frac{\partial}{\partial x_i} \cdot (f_j \underline{e}_j) = \underline{e}_i \cdot \underline{e}_j \frac{\partial f_j}{\partial x_i} = \frac{\partial f_j}{\partial x_i}$$

same idea

→ Also, whenever you see a term like $\frac{\partial f_k}{\partial x_i}$, you now know $\frac{\partial f_k}{\partial x_i} = \nabla \cdot \underline{f}$.

c. An identity using index notation: let $\phi(x)$ be a scalar field

$$\nabla \cdot (\phi \underline{f}) = \underline{e}_i \frac{\partial}{\partial x_i} \cdot (\phi f_j \underline{e}_j)$$

(i) use product rule
(ii) \underline{e}_j are constant vectors
(iii) the inner product (\cdot)

$$= (\underline{e}_i \cdot \underline{e}_j) \frac{\partial}{\partial x_i} (\phi f_j)$$

only operates on vectors, not the scalar components

$$= \delta_{ij} \left[\frac{\partial \phi}{\partial x_i} f_j + \phi \frac{\partial f_j}{\partial x_i} \right]$$

$$= \frac{\partial \phi}{\partial x_j} f_j + \phi \frac{\partial f_j}{\partial x_j} = (\nabla \phi) \cdot \underline{f} + \phi \nabla \cdot \underline{f}$$

$$\therefore \nabla \cdot (\phi \underline{f}) = (\nabla \phi) \cdot \underline{f} + \phi \nabla \cdot \underline{f}$$

← notice how similar this is to the normal product rule of differentiation
(134)

Exercise: Evaluate (i) $\nabla \cdot (\underline{a} \wedge \underline{b})$

(ii) $\nabla (\underline{a} \cdot \underline{b})$

where $a(x)$ and $b(x)$ are vector functions of position

7. Curl of a vector field: $\underline{\nabla}^1 f$ or sometimes $\text{curl } \underline{f}$

a. Let us first calculate using standard ideas

$$\underline{\nabla}^1 f = e_i \frac{\partial}{\partial x_i} \wedge (f_j e_j)$$

As before, the e_j are constant vectors. Also, the $\text{curl}(\wedge)$ operation only affects vector

$$= (e_i \wedge e_j) \frac{\partial f_j}{\partial x_i}$$

c. $\underline{\nabla}^1 f = \epsilon_{ijk} \frac{\partial f_j}{\partial x_i} e_k$ (135)

(remember: the summation convention is in effect. Write out what eqn 135 represents.)

NOTE: Sometimes you will find that people write

$$(\underline{\nabla}^1 f)_k = \epsilon_{ijk} \frac{\partial f_j}{\partial x_i}$$

\uparrow indicates the k^{th} component of the vector $\underline{\nabla}^1 f$.

b. Also, since $\epsilon_{ijk} = \epsilon_{kij}$ we may write

$$\underline{\nabla}^1 f = \epsilon_{kij} \frac{\partial f_j}{\partial x_i} e_k$$

c. Alternatively, let's simply show that eqn (135) agrees with what you have seen in your earlier vector calculus course.

Since

$\underline{\nabla}^1 f = (e_i \wedge e_j) \frac{\partial f_j}{\partial x_i}$ and since the summation convention has been assumed and the variables i, j appear twice, we must sum $i=1 \rightarrow 3, j=1 \rightarrow 3$.

then

$$\underline{\nabla}^1 f = (e_1 \wedge e_1) \frac{\partial f_1}{\partial x_1} + (e_1 \wedge e_2) \frac{\partial f_2}{\partial x_1} + (e_1 \wedge e_3) \frac{\partial f_3}{\partial x_1} + (e_2 \wedge e_1) \frac{\partial f_1}{\partial x_2} + (e_2 \wedge e_2) \frac{\partial f_2}{\partial x_2} + (e_2 \wedge e_3) \frac{\partial f_3}{\partial x_2}$$

$$= e_1 \left(\frac{\partial f_3}{\partial x_2} - \frac{\partial f_2}{\partial x_3} \right) + e_2 \left(\frac{\partial f_1}{\partial x_3} - \frac{\partial f_3}{\partial x_1} \right) + e_3 \left(\frac{\partial f_1}{\partial x_2} - \frac{\partial f_2}{\partial x_1} \right)$$

$$= \begin{vmatrix} e_1 & e_2 & e_3 \\ \frac{\partial}{\partial x_1} & \frac{\partial}{\partial x_2} & \frac{\partial}{\partial x_3} \\ f_1 & f_2 & f_3 \end{vmatrix}$$

which is probably how you previously saw it represented.

8. Two examples

a. An important identity: Evaluate $\nabla \wedge \nabla \phi(x)$

Using index notation we have

$$\begin{aligned}\nabla \wedge \nabla \phi &= e_i \frac{\partial}{\partial x_i} \wedge (e_j \frac{\partial \phi}{\partial x_j}) = e_i \wedge e_j \frac{\partial^2 \phi}{\partial x_i \partial x_j} \\ &= \epsilon_{ijk} \frac{\partial^2 \phi}{\partial x_i \partial x_j} e_k\end{aligned}$$

Next, notice that by using the 'indices-interchange' property of ϵ_{ijk} we may write

$$\nabla \wedge \nabla \phi = \epsilon_{ijk} \frac{\partial^2 \phi}{\partial x_i \partial x_j} e_k = -\epsilon_{jik} \frac{\partial^2 \phi}{\partial x_i \partial x_j} e_k = -\epsilon_{jik} \frac{\partial^2 \phi}{\partial x_j \partial x_i} e_k$$

↑
relabeled dummy variables: $i \rightarrow j$
 $j \rightarrow i$

By comparing the first and last equalities:

$$\epsilon_{ijk} \frac{\partial^2 \phi}{\partial x_i \partial x_k} = -\epsilon_{ijk} \frac{\partial^2 \phi}{\partial x_k \partial x_i} = 0 \quad (\text{provided } \frac{\partial^2 \phi}{\partial x_i \partial x_k} = \frac{\partial^2 \phi}{\partial x_k \partial x_i})$$

Exercise:
 $\underline{g} \wedge \underline{g} = 0$, then
 $g_i g_i \epsilon_{ijk} e_k = 0$
 You should recognize
 this combination is $= 0$.

∴

$$\nabla \wedge \nabla \phi = 0 \quad \text{for any continuously differentiable scalar function } \phi(x).$$

b. Consider two vectors $\underline{a}(x)$ and $\underline{b}(x)$.

⇒ Evaluate $\nabla \cdot (\underline{a} \wedge \underline{b})$

SOLN

$$e_i \frac{\partial}{\partial x_i} \cdot (a_j e_j \wedge b_k e_k) = e_i \frac{\partial}{\partial x_i} (a_j b_k) \epsilon_{jkl} e_l$$

constant, so may be taken outside of parentheses

$$= (\epsilon_i \cdot e_l) \frac{\partial}{\partial x_i} (a_j b_k) \epsilon_{jkl} = \frac{\partial}{\partial x_i} (a_j b_k) \epsilon_{jkl} \quad \epsilon_{jkl} = \epsilon_{ijk}$$

$$= (\frac{\partial a_j}{\partial x_i}) \epsilon_{ijk} b_k + \frac{\partial b_k}{\partial x_i} \epsilon_{kij} a_j$$

$$\nabla \cdot (\underline{a} \wedge \underline{b}) = \frac{\partial a_j}{\partial x_i} \epsilon_{ijk} b_k - \frac{\partial b_k}{\partial x_i} \epsilon_{ijk} a_j = (\nabla \cdot \underline{a}) b_k - (\nabla \cdot \underline{b}) a_j$$

or in vector notation,


$$\nabla \cdot (\underline{a} \wedge \underline{b}) = (\nabla \cdot \underline{a}) \cdot \underline{b} - (\nabla \cdot \underline{b}) \cdot \underline{a}$$

9. A menagerie of typical calculations

a. In many problems a scalar or vector function, $\phi(\underline{x})$ or $\underline{f}(\underline{x})$ say, depends on position \underline{x} .

let \underline{x} = position vector

let $r^2 = \underline{x} \cdot \underline{x} = x_i^2$ denote distance from the origin.



b. What is ∇r^2 ?

$$\nabla r^2 = \sum_j \frac{\partial}{\partial x_j} r^2 = \sum_j 2r \frac{\partial r}{\partial x_j}. \quad (*)$$

what is this?

... continue in a moment, but first let's address what is ∇r or in component form $\frac{\partial r}{\partial x_j} = ?$

c. Since $r = \sqrt{x_1^2 + x_2^2 + x_3^2} = \sqrt{x_1^2 + x_2^2 + x_3^2}$ we see that

$$\frac{\partial r}{\partial x_1} = \frac{x_1}{\sqrt{x_1^2 + x_2^2 + x_3^2}} = \frac{x_1}{r} \quad ; \text{ similarly } \frac{\partial r}{\partial x_2} = \frac{x_2}{r}; \quad \frac{\partial r}{\partial x_3} = \frac{x_3}{r}$$

$$\text{and so } \frac{\partial r}{\partial x_j} = \frac{x_j}{r} \quad \text{or} \quad \nabla r = \sum_j \frac{\partial r}{\partial x_j} = \sum_j \frac{x_j}{r} = \frac{\underline{x}}{r}$$

$$\therefore \boxed{\frac{\partial r}{\partial x_j} = \frac{x_j}{r}}_{(136a)}; \quad \boxed{\nabla r = \frac{\underline{x}}{r}}_{(136b)}$$

Alternatively, we may note that $r^2 = x_i x_i$
so taking $\frac{\partial}{\partial x_j}$ gives

$$2r \frac{\partial r}{\partial x_j} = \frac{\partial}{\partial x_j} (x_i x_i) = 2 x_i \frac{\partial x_i}{\partial x_j} = 2 x_j$$

$$\therefore \boxed{\frac{\partial r}{\partial x_j} = \frac{x_j}{r}}$$

(note the simplicity of the index notation calculation.)

d. ... and so now we return to $(*)$ to see that $\nabla r^2 = \sum_j 2r \cdot \frac{x_j}{r} = 2 \underline{x}$

e. Evaluate $\nabla \cdot (\underline{x} e^r) = (\nabla \cdot \underline{x}) e^r + \underline{x} \cdot \nabla e^r = 3e^r + x_i \frac{\partial}{\partial x_i} e^r \rightarrow \text{use chain rule + results above}$
 $= 3e^r + x_i \frac{\partial r}{\partial x_i} \frac{\partial}{\partial r} e^r = 3e^r + x_i \cdot \frac{x_i}{r} e^r = (3+r)e^r.$

f. Exercise: evaluate $\nabla^2 \underline{x}$.

D. Integral Theorems

1. DIVERGENCE THEOREM (also, sometimes referred to as Gauss's Theorem, Green's or Ostrogradsky's Theorem)

⇒ This theorem relates integrals over volumes to integrals over the bounding surfaces.

a. The Divergence Theorem states that given a continuous vector function $\underline{f}(x)$ with continuous first partial derivatives, then

$$\text{DIVERGENCE THEOREM} \quad \int_V \nabla \cdot \underline{f} \, dV = \int_S \underline{n} \cdot \underline{f} \, dS \quad (137)$$

volume → V \underline{n} denotes ALL bounding surfaces S

where \underline{n} is the unit outward normal from V



b. You may wish to read & recall the proof of the theorem given in Math 21. We will provide a proof for the two dimensional situation.

c. Eqn (137), written out in 3D, is

$$\int_V \left(\frac{\partial f_1}{\partial x_1} + \frac{\partial f_2}{\partial x_2} + \frac{\partial f_3}{\partial x_3} \right) dV = \int_S (n_1 f_1 + n_2 f_2 + n_3 f_3) dS$$

or using index notation

$$\int_V \frac{\partial f_i}{\partial x_i} dV = \int_S n_i f_i dS$$

notation:
Sometimes see
 $dS = \underline{n} d$

① When writing this theorem in vector form, it is useful to get into the habit of writing \underline{n} on the left. The order will be important when we apply the Divergence Theorem to tensor functions.

② It is common to see the volume integral indicated as \iiint_V and the surface integral as \iint_S . I will always use the simple notation shown in eqn (136) and clearly indicate volume or surface integration with V, dV or S, dS respectively.

(2D)

2. Planar^{^*} versions of the Divergence Theorem


(common to discuss
these "introductory
vector calculus courses.)

- a. Consider some area A in the plane bounded by a curve C .

The Divergence Theorem (eqn 13) is now

$$(136a) \int_A \nabla \cdot \underline{f} dA = \oint_C \underline{n} \cdot \underline{f} ds$$

$C \leftarrow$ integral around boundary



- b. You have probably seen the Divergence Theorem in a somewhat different form.

Let \underline{n} and \underline{t} denote the unit outward normal and tangent vectors along the boundary C (see sketch).

Let $d\underline{x}$ indicate a small displacement along the boundary ($ds = |d\underline{x}|$)

$$\text{Since } \underline{t} = \frac{d\underline{x}}{ds} \Rightarrow \underline{t} ds = dx_1 \underline{e}_1 + dx_2 \underline{e}_2$$

Also, as $\underline{n} \cdot \underline{t} = 0$, then

$$\underline{n} = \frac{dx_2 \underline{e}_1}{ds} - \frac{dx_1 \underline{e}_2}{ds} \rightarrow \underline{n} ds = dx_2 \underline{e}_1 - dx_1 \underline{e}_2$$

- c. If we simply expand eqn (136a) then

$$\int_A \left(\frac{\partial f_1}{\partial x_1} + \frac{\partial f_2}{\partial x_2} \right) dA = \oint_C \underline{n} \cdot \underline{f} ds = \oint_C (f_1 dx_2 - f_2 dx_1) \quad (136)$$

which is a common eqn to see.

Alternatively, if we let $f_1(x_1, x_2) = N(x_1, x_2)$
 $f_2(x_1, x_2) = -M(x_1, x_2)$ then

then we may write

$$dx_1 dx_2 = dx dy$$

$$\int_A \left(\frac{\partial N}{\partial x_1} - \frac{\partial M}{\partial x_2} \right) d\overline{A} = \oint_C (M dx_1 + N dx_2) \quad (136c)$$

which also commonly appears in introductory vector calculus courses.

d. Proof of the Divergence Theorem in 2D


a. A result we will need: $\int_{x=a}^b \frac{\partial g}{\partial x}(x, y) dx = g(b, y) - g(a, y)$. *Integration of a function with respect to x, y*

b. Consider $\int_A \left(\frac{\partial f_x}{\partial x} + \frac{\partial f_y}{\partial y} \right) dA$

Let the region A be such that any vertical line or horizontal line intersects the boundary C in at most two points (see sketch)

→ identify functions $x_e(y), x_h(y)$ which describe the boundary as the y -direction is traversed from c to d .

→ Similarly, $y_e(x), y_h(x)$ describe the boundary as the x -direction is traversed from a to b .



c. Then,

$$\begin{aligned} \int_A \nabla \cdot \underline{f} dA &= \int_{y=c}^d \int_{x=x_e(y)}^{x_h(y)} \frac{\partial f_x}{\partial x} dx dy + \int_{x=a}^b \int_{y=y_e(x)}^{y_h(x)} \frac{\partial f_y}{\partial y} dy dx \\ &= \int_{y=c}^d [f_x(x_h, y) - f_x(x_e, y)] dy + \int_{x=a}^b [f_y(x, y_h) - f_y(x, y_e)] dy \quad \text{note the form of t} \\ &= \int_{y=c}^d f_x(x_h(y), y) dy + \int_d^c f_x(x_e(y), y) dy - \int_{x=a}^b f_y(x, y_h(x)) dy - \int_a^x f_y(x, y_e(x)) dx \\ &= \oint_C f_x(x, y) dy - \oint_C f_y(x, y) dx \end{aligned}$$

$$\therefore \int_A \nabla \cdot \underline{f} dy = \oint_C (f_x dy - f_y dx) = \oint_C \underline{n} \cdot \underline{f} ds$$

proof complete

Remark: It is straightforward to extend the theorem to more general regions by subdividing the region into smaller regions similar to the figure above.

where \oint denotes integration in the counter-clockwise direction and $\underline{n} = (\frac{dy}{ds}, -\frac{dx}{ds})$.

Note: It is not terribly useful to attempt to memorize all the different versions in which the Divergence Theorem may be written. Rather simply remember

$$\int_V \underline{\nabla} \cdot \underline{f} dV = \int_S \underline{n} \cdot \underline{f} dS \text{ in 3D} \quad \text{or} \quad \int_A \underline{\nabla} \cdot \underline{f} dA = \int_C \underline{n} \cdot \underline{f} ds \text{ in 2D}$$

and recognize that with a little algebra & geometry you can derive most intermediate formulas.

c. A planar version of Stokes' Theorem

Recall eqn (13E) on the bottom of pg. 141.

Identify the scalar function $N(x_1, x_2)$ and $M(x_1, x_2)$ with the components of a vector according to

$$\underline{q}(x) = q_1(x_1, x_2) \underline{e}_1 + q_2(x_1, x_2) \underline{e}_2 = M(x_1, x_2) \underline{e}_1 + N(x_1, x_2) \underline{e}_2$$

Then, from (13E) we have

$$\int_A \left(\frac{\partial q_2}{\partial x_1} - \frac{\partial q_1}{\partial x_2} \right) dA = \oint_C q_1 dx_1 + q_2 dx_2$$

which we may express in a simple form using vector notation

$$(13g) \quad \boxed{\int_A (\underline{\nabla} \times \underline{q}) \cdot \underline{e}_3 dA = \oint_C \underline{q} \cdot \underline{t} ds}$$

where $\underline{t} = \frac{dx_1}{ds} \underline{e}_1 + \frac{dx_2}{ds} \underline{e}_2$
 $= \frac{d\underline{x}}{ds}$ b the unit tangent vector

↑ Planar version of Stokes Theorem.

'we'll come back to this in a few pages.'

3. Some theorems, or identities, which follow directly from the Divergence Theorem

Begin with $\int_V \nabla \cdot \underline{f} dV = \int_S \underline{n} \cdot \underline{f} dS \quad (*)$

or you may wish to write $\int_V \frac{\partial f_i}{\partial x_i} dV = \int_S n_i f_i dS$

• \rightarrow In the calculation presented below I will work in vector notation. You may wish instead to work in index notation.

a. Let $\underline{f} = \phi \underline{b}$ where $\phi = \phi(\underline{x})$ and \underline{b} is an arbitrary constant vec

Substitute into (*).

Since \underline{b} = constant vector, $\nabla \cdot (\phi \underline{b}) = \underline{b} \cdot \nabla \phi$
So, we arrive at

(**) $\underline{b} \cdot \int_V \nabla \phi dV = \underline{b} \cdot \int_S \underline{n} \phi dS$

where \underline{b} can be taken outside of the integrals because it is a constant vector.

Because this eqn must be true for arbitrary \underline{b} , we conclude

$$\int_V \nabla \phi dV = \int_S \underline{n} \phi dS \quad (140)$$

which you may like to refer to as the 'Divergence' Theorem for scalar functions.

We will make use of this argument several times in this course.

c. The formal argument to go from (**) to (140) goes like this:

Rearrange (*) as

(141) $\underline{b} \cdot \left[\int_V \nabla \phi dV - \int_S \underline{n} \phi dS \right] = 0$

where the quantity inside the brackets [] is clearly a vector.

Remember \underline{b} has been assumed to be an arbitrary constant vector

Thus from (141) we must either have

- (i) $[\underline{b}] = 0$
- (ii) $\underline{b} = 0$
- (iii) $\underline{b} \perp [\underline{b}]$

} but \underline{b} is arbitrary so (ii) & (iii) cannot hold.
Therefore $[\underline{b}] = 0$

$\therefore \int_V \nabla \phi dV = \int_S \underline{n} \phi dS$

or in index notation $\int \frac{\partial \phi}{\partial x_i} e_i = \int_S n_i \phi dS$

d. Another theorem which follows from the Divergence Theorem is obtained by letting

$$\underline{f} = \underline{\nabla} \phi$$

$$\begin{aligned} \text{Then since } \underline{\nabla} \cdot \underline{f} &= \underline{\nabla} \cdot (\underline{\nabla} \phi) = \sum_i \frac{\partial}{\partial x_i} \cdot \left(\sum_j \frac{\partial \phi}{\partial x_j} \right) \\ &= \frac{\partial^2 \phi}{\partial x_i \partial x_i} = \frac{\partial^2 \phi}{\partial x_i^2} = \frac{\partial^2 \phi}{\partial x_1^2} + \frac{\partial^2 \phi}{\partial x_2^2} + \frac{\partial^2 \phi}{\partial x_3^2} = \nabla^2 \phi \end{aligned}$$

we have

the LAPLACIAN

$$\int_V \nabla^2 \phi \, dV = \int_S \underline{n} \cdot \underline{\nabla} \phi \, dS \quad (141)$$

$\frac{\partial \phi}{\partial n}$ ← gradient of ϕ in the direction of the unit normal \underline{n} (i.e., directional derivative along \underline{n})

4. GREEN'S THEOREMS (Hildebrand, p 301-2; Kreysig §9.8)

a. Begin with the Divergence Theorem and let $\underline{f} = \Psi \underline{\nabla} \phi$.

Then,

$$\begin{aligned} \int_S (\Psi \underline{n} \cdot \underline{\nabla} \phi) \, dS &= \int_V \underline{\nabla} \cdot (\Psi \underline{\nabla} \phi) \, dV = \int_V \sum_i \frac{\partial}{\partial x_i} \cdot \left(\Psi \frac{\partial \phi}{\partial x_i} \underline{e}_j \right) \\ &= \int_V \frac{\partial}{\partial x_i} \left(\Psi \frac{\partial \phi}{\partial x_i} \right) \, dV = \int_V \left(\frac{\partial \Psi}{\partial x_i} \frac{\partial \phi}{\partial x_i} + \Psi \frac{\partial^2 \phi}{\partial x_i^2} \right) \, dV \end{aligned}$$

$$\therefore \int_S \Psi \underbrace{\underline{n} \cdot \underline{\nabla} \phi}_{\frac{\partial \phi}{\partial n}} \, dS = \int_V [\Psi \underline{\nabla} \cdot \underline{\nabla} \phi + \Psi \nabla^2 \phi] \, dV \quad (142)$$

Green's
1st form
or Green's 1st
Identity

b. Interchange Ψ, ϕ in the above eqn, then subtract from the eqn just derived.

$$\Rightarrow \int_S \left(\Psi \frac{\partial \phi}{\partial n} - \phi \frac{\partial \Psi}{\partial n} \right) \, dS = \int_V (\Psi \nabla^2 \phi - \phi \nabla^2 \Psi) \, dV \quad (143)$$

Green's
2nd form
or
Green's Secnd
Identity

Exercise : derive this eqn

An interesting aside: We will return to this idea when we study pdes. Just skim for now.

14c

Green's Theorems are often useful for proving some very general results.

For example, begin with Green's first form; let $\Psi = \phi$

Then,

$$\int_S \phi \frac{\partial \phi}{\partial n} dS = \int_V [\nabla \phi \cdot \nabla \phi + \phi \nabla^2 \phi] dV$$

p.d.e

Now suppose you wish to solve $\nabla^2 \phi = 0$ (Laplace's eqn) in V subject to the b.c. $\phi = 0$ on S

What is $\phi(x)$ for $x \in V$?

$\phi(x) = ?$ S $\rightarrow V$
 $\phi = 0$

Well, we are given $\phi = 0$ on the boundary S so,

$$\int_S \phi \frac{\partial \phi}{\partial n} dS = 0.$$

Furthermore, since $\nabla^2 \phi = 0$ in V , the eqn above reduces to

$$\int_V [\nabla \phi \cdot \nabla \phi] dV = 0$$

But the integrand $\nabla \phi \cdot \nabla \phi$ is the sum of squares, so is always positive. For the integral to vanish we must consequently require $\nabla \phi \cdot \nabla \phi = 0$ everywhere in V

or $\nabla \phi = 0$, i.e., ϕ doesn't vary with spatial position in V
 $\Rightarrow \phi = \text{constant}$ in V

But $\phi = 0$ on S and so therefore $\phi = 0$ throughout V

#

5. A further generalization of the Divergence Theorems

a. We began by stating the Divergence Theorem

Where S represents the closed surface enclosing the volume V and \underline{n} is the unit outward normal from the volume.

$$\int_V \nabla \cdot \underline{f} dV = \int_S \underline{n} \cdot \underline{f} dS$$

We then proved a form suitable for scalar functions,

$$\int_V \nabla \phi dV = \int_S \underline{n} \cdot \underline{\phi} dS$$

\leftarrow eqn (140), p. 144.

or using index notation

$$\int_V \frac{\partial \phi}{\partial x_i} dx_i = \int_S n_i \phi dS \quad \epsilon_i$$

If two vectors are equal, their corresponding components are equal, so we see that

$$(*) \quad \int_V \frac{\partial \phi}{\partial x_i} dx_i = \int_S n_i \phi dS \quad \text{for } i=1, 2 \text{ or } 3.$$

b. We now construct a form useful for when the cross-product appears.

For example, consider integrals of the form $\int_V \nabla^1 f dV$

It is simplest to work in index notation. Notice that $\nabla^1 f$ represents a vector so the result of the integration is also a vector. Basically, we then proceed by considering each component of the vector separately.

So using index notation

$$\int_V \nabla^1 f dV = \int_V \frac{\partial f_k}{\partial x_j} \epsilon_{jki} \epsilon_i dV = \int_V \frac{\partial f_k}{\partial x_j} dV \underbrace{\epsilon_{jki} \epsilon_i}_{\text{these are just constants}}$$

But for each k and i we know that

$$\int_V \frac{\partial f_k}{\partial x_j} dV = \int_S n_j f_k dS$$

(remember - the summation convention is in effect so this actually represents the sum of lots of terms)

In other words, for each k, i use eqn (*) and let $f_k = \phi$ so that

$$\int_V \frac{\partial \phi}{\partial x_j} dV \epsilon_{jki} \epsilon_i = \int_S n_j \phi dS \epsilon_{jki} \epsilon_i$$

Hence,

$$\int_V \nabla^1 f dV = \int_V \frac{\partial f_k}{\partial x_j} dV \epsilon_{jki} \epsilon_i = \int_S n_j f_k dS \epsilon_{jki} \epsilon_i = \int_S \underline{n}^1 f dS \quad (144)$$

C. Notice, that we can conveniently summarize all of the above results concerning the Divergence Theorem, as follows:

$$\int_V \nabla \cdot \vec{F} \, dV = \int_S \vec{n} \cdot \vec{F} \, ds \quad (145)$$

where Φ is any quantity, scalar or vector, and $*$ is any operation (scalar product, vector product or a simple gradient operation) that makes sense.

d. EXAMPLES :

(i) let $\underline{a} = \text{constant vector}$

$$\int_S \underline{n} \cdot \underline{g} \, dS = \int_V \underline{g} \cdot \underline{a} \, dV = 0$$

V 0 if $\underline{g} = \text{curl}$
vector

(ii) Evaluate

$$\int \underline{n} \cdot (\nabla^{\wedge} f) \, ds$$

By the ⁵ Divergence Theorem,

$$\int_S \underline{n} \cdot (\nabla^V f) dS = \int_V \underline{\nabla} \cdot (\nabla^V f) dV = 0.$$

(ii) Evaluate $\int_S \mathbf{n} \cdot \nabla r^2 dS$

Using index notation, $\nabla \cdot \vec{r}^2 = \eta \cdot \frac{\partial}{\partial x_i} (r^2) \rightarrow \frac{\partial}{\partial x_i} (r^2) = 2r \frac{\partial r}{\partial x_i} = 2x_i$

$$\text{So, } \int_S n \cdot \nabla r^2 dS = \int_S n_i \frac{\partial}{\partial x_i} (r^2) dS \quad \text{since } \frac{\partial r}{\partial x_i} = x_i/r \text{ (pg. 139)}$$

$$= \int_S 2 n_i x_i dS = 2 \int_V \frac{\partial x_i}{\partial x_i} dV \text{ by the Divergence Theorem}$$

$$= 2 \cdot \delta_{ii} \int_V dV = V \text{ where } V \text{ is volume of domain; } \delta_{ii} =$$

$\int_S n \cdot \nabla r^2 dS = 6V$ where V is volume of domain bounded.

C. Notice, that we can conveniently summarize all of the above results concerning the Divergence Theorem, as follows:

$$\int_V \nabla * \Phi \, dV = \int_S n * \Phi \, dS \quad (145)$$

where Φ is any quantity, scalar or vector, and $*$ is any operation (scalar product, vector product or a simple gradient operation) that makes sense.

d. EXAMPLES :

(i) let \underline{a} = constant vector $\int_S n \cdot \underline{a} \, dS = \int_V \underline{a} \cdot \underline{a} \, dV = 0$
 $V \quad 0 \text{ if } \underline{a} = \text{constant vector}$

(ii) Evaluate $\int_S n \cdot (\nabla^1 f) \, dS$

By the Divergence Theorem,

$$\int_S n \cdot (\nabla^1 f) \, dS = \int_V \nabla \cdot (\nabla^1 f) \, dV = 0.$$

0 for any twice continuously differentiable vector function f .

(iii) Evaluate $\int_S n \cdot \nabla r^2 \, dS$

Using index notation, $n_i \nabla r^2 = n_i \frac{\partial}{\partial x_i} (r^2) \rightarrow \frac{\partial}{\partial x_i} (r^2) = 2r \frac{\partial r}{\partial x_i} = 2x_i$

so, $\int_S n_i \nabla r^2 \, dS = \int_S n_i \frac{\partial}{\partial x_i} (r^2) \, dS \quad \text{since } \frac{\partial r}{\partial x_i} = x_i/r \text{ (pg. 139)}$

$$= \int_S 2 n_i x_i \, dS = 2 \int_V \frac{\partial x_i}{\partial x_i} \, dV \text{ by the Divergence Theorem}$$

$$= 2 \cdot \delta_{ii} \int_V \, dV = V \text{ volume of domain; } \delta_{ii} = 1$$


$\therefore \boxed{\int_S n \cdot \nabla r^2 \, dS = 6V}$ where $V \equiv \text{volume of domain}, \text{ bounded by } \Sigma$

e. For those of you who would like more exercises, show

$$(i) \int_S \underline{n} \cdot \nabla r^2 dS = 0$$

$$(ii) \int_S \underline{n} \cdot \nabla (\underline{x} \cdot \underline{a}) dS = 0 \quad \text{where } \underline{a} \text{ is a constant vector and } \underline{x} \text{ is a position vector to a pt on the surface}$$

$$r = |\underline{x}|$$




6. There is one final important point concerning the Divergence Theorem, and that is if there are multiple bounding surfaces, you must include all of them.

In other words

$$\int_V \nabla \cdot \underline{f} dV = \int_S \underline{n} \cdot \underline{f} dS$$

\nwarrow all bounding surfaces

So, if V is as shown below:



Then

$$\int_S \underline{n} \cdot \underline{f} dS = \int_{S_1 + S_2 + S_3 + S_4} \underline{n} \cdot \underline{f} dS$$


and notice how on each surface, \underline{n} points outward from V

7. Stokes' Theorem - This allows one to express an integral around a closed curve as an integral over that area having the curve as a boundary.

Q. Let C be a closed curve and let S be a surface which has C as a bounding edge. (imagine a 'hat-shaped' surface)

let \underline{n} = unit normal to S with direction given by the right-hand rule - as you curl your hand in the direction indicated about C , your thumb points in the direction of \underline{n} .

\underline{t} = unit tangent vector to C .



sometimes people write $ds = \underline{t} ds$

Stokes Theorem :

$$\oint_C f \cdot \underline{t} \, ds = \int_S \underline{n} \cdot (\nabla f) \, dS \quad (H6)$$

Recall proof given in Math 21.

↑
differential
element along
 C

← distinguish
'little s'
and
'big S'

Using index notation :

$$\oint_C f_i t_i \, ds = \int_S n_i \frac{\partial f_k}{\partial x_j} \epsilon_{ijk} \, dS \quad (H7)$$

Writing this all out

$$\oint_C (f_1 t_1 + f_2 t_2 + f_3 t_3) \, ds = \int_S \left[n_1 \left(\frac{\partial f_3}{\partial x_2} - \frac{\partial f_2}{\partial x_3} \right) + n_2 \left(\frac{\partial f_1}{\partial x_3} - \frac{\partial f_3}{\partial x_1} \right) + n_3 \left(\frac{\partial f_2}{\partial x_1} - \frac{\partial f_1}{\partial x_2} \right) \right] dS$$

NOTE: This is true for an arbitrary surface S with C as a bounding edge.

b. We can actually construct a simple proof using results developed so far


Using s to denote arclength along the bounding curve C ,

$$\frac{ds}{ds} = \underline{t} \text{ - then } dx_i = t_i \, ds$$

So, considering the f_1 term :

$$\oint_C f_1 t_1 \, ds = \oint_C f_1 \, dx_1 = \oint_{C'} f(x_1, x_3) \, dx_1$$

function of x_1, y, z $\rightarrow z = z(x_1, y)$ represents curve C given C' in xy -plane



and we have written
 $f_1(x_1, x_2, x_3(x_1, x_2)) = f(x_1, x_2)$

where we have essentially projected information down to the

b. "simple proof" (continued)

With information "in the plane" we can now use the identity given on pg. 147:

one form of the 'Divergence Theorem' was $\int_S n_i \phi dS = \int_V \frac{\partial \phi}{\partial x_i} dV$
which has the 'planar version'

$$\int_A \frac{\partial \phi}{\partial x_i} dA = \int_C \phi n_i ds$$

or

$$\int_A \frac{\partial \phi}{\partial x_1} dA = \int_C \phi dx_2 \quad \text{and} \quad \int_A \frac{\partial \phi}{\partial x_2} dA = - \int_C \phi dx_1$$

$\int_C t ds = dx_1 \varepsilon_1 + dx_2 \varepsilon_2$
 $n ds = dx_2 \varepsilon_1 - dx_1 \varepsilon_2$

Hence, beginning with the eqn on the bottom of the last page

$$\begin{aligned} \int_C \phi f_1 dx_1 &= - \int_A \frac{\partial f_1}{\partial x_2} dA \quad \text{but } \frac{\partial}{\partial x_2} f_1(x_1, x_2) = \frac{\partial}{\partial x_2} f_1(x_1, x_2, x_3(x_1, x_2)) \\ &= - \int_A \left[\frac{\partial f_1}{\partial x_2} + \frac{\partial x_3}{\partial x_2} \frac{\partial f_1}{\partial x_3} \right] dA \quad \text{by the chain-rule} \end{aligned}$$

Now, a differential element dS in space is related to its projection

in the xy -plane by $dA = n_3 dS$

and furthermore one can show

i.e.,

$$\frac{\partial x_3}{\partial x_2} = - \frac{n_2}{n_3}$$

so that we have

$$\int_C \phi f_1 dx_1 = - \int_S \left[\frac{\partial f_1}{\partial x_2} n_3 - \frac{\partial f_1}{\partial x_3} n_2 \right] dS$$

or

$$\int_C \phi f_1 dx_1 = \int_C \phi f_1 t_1 ds = \int_S \left(\frac{\partial f_1}{\partial x_3} n_2 - \frac{\partial f_1}{\partial x_2} n_3 \right) dS \leftarrow$$

which accounts for two of the terms in Stokes theorem. In a similar manner one can account for the other 2 terms. As in standard versions of these proofs, it is necessary to imagine that S is subdivided into sections which can be projected onto the xy -plane, even when the whole of S does not have a one-to-one projection.

