

ECE609 Spring07  
**HOMEWORK 1**  
Review of Quantum Mechanics

### 1 The Photoelectric effect (15pts)

1. A metal has a workfunction of 4.3 eV. What is the minimum photon energy in Joule to emit an electron from this metal through the photo-electric effect? What are the photon frequency in Terahertz and the photon wavelength in micrometer (we note that  $\lambda = c/\nu$  where  $c$  is the velocity of the light  $\simeq 3 * 10^8 m.s^{-1}$ )? What is the corresponding photon momentum ?
2. A photocathode in Potassium has a workfunction of 2 eV. What condition for the photon wavelength is necessary (in nanometer) to allow the photoelectric effect ?, For an incoming radiation of  $\lambda = 546.1nm$ , what is the maximum velocity of an extracted electron ? For an incoming radiation of  $\lambda = 645.5nm$ , what is the maximum velocity of an extracted electron ?

### 2 Hydrogen atom (30pts)

The fundamental level (1s state) of the hydrogen atom can be described by the wave function (**spherical coordinates**  $r \geq 0$ ):

$$\Psi(r) = Cexp(-r/a_0),$$

where  $C$  is a constant (complex) and  $a_0$  is defined as the Bohr radius.

(Hint: for all the integrations, we will consider a volume  $4\pi r^2 dr$  - elementary volume between two spheres with radius  $r$  and  $r + dr$  (shell))

1. Find the value of  $C$ ,
2. What is the density of probability to find an electron in a volume  $4\pi r^2 dr$  (shell). Plot the density of probability  $dP/dr$  in function of  $r$ .
3. We call  $r = r_p$  the value for which this probability exhibits a maximum. What is  $r_p$  in function of  $a_0$ ?  
Comment
4. What is the probability to find an electron at the distance between  $0.9r_p$  and  $1.1r_p$  ?
5. Give the average value (expectation value) of the distance that separate the proton from the electron  $\langle r \rangle$ . We note  $\langle r \rangle = \langle \Psi | r | \Psi \rangle = \int_0^\infty \Psi^*(x)r\Psi(x)4\pi r^2 dr$ .
6. Give the expectation values (in eV) of the potential energy  $\langle U \rangle$  and the kinetic energy  $\langle T \rangle$ . What is the value of the total energy ? Comment.

### 3 Finite potential well (25pts)

We consider a 1D finite potential well, where the variations of  $U(x)$  are defined by:

$$U(x) = \begin{cases} 0 & \text{if } x < -a/2 \text{ (region I), or } x > a/2 \text{ (region III)} \\ -U_0 & \text{if } -a/2 < x < a/2 \text{ (region II)} \end{cases}$$

where  $a$  and  $U_0$  are positive constants. We consider an incoming electron wave from the left of the system with energy  $E > 0$  and amplitude equal to one.

1. Give the expressions of the wave function  $\Psi(x)$  in the three regions. We will set:

$$k = (2mE)^{1/2}/\hbar \text{ and } K = (2m(E + U_0))^{1/2}/\hbar.$$

2. The Transmission coefficient  $T$  can take the following form:

$$T = \frac{1}{1 + f(E)\sin^2 g(E)}$$

where

$$f(E) = \frac{U_0^2}{4E(E + U_0)}$$

and

$$g(E) = (2m(E + U_0))^{1/2}a/\hbar$$

What is the reflection coefficient  $R$  (no calculation is necessary here)?

3. Numerical application,  $a = 1.5A$ ,  $U_0 = 16eV$ . Plot  $T(E)$  for  $E < 150eV$
4. Calculate (analytically) the energy values for which  $T$  is equal to 1 (total transmission). What are the numerical expressions of these energy values (we consider only the values smaller than  $150eV$ ) ?
5. What is happening if  $E \rightarrow \infty$ . Comment.

## 4 Finite potential wall (30pts)

We consider a 1D finite potential wall, where the variations of  $U(x)$  are defined by:

$$U(x) = \begin{cases} 0 & \text{if } x \leq 0 \text{ (region I), or } x \geq a \text{ (region III)} \\ U_0 & \text{if } 0 < x < a \text{ (region II)} \end{cases}$$

where  $a$  and  $U_0$  are positive constants.

We consider an incoming electron wave from the left of the system with energy  $0 < E < U_0$  and amplitude equal to one.

1. Show that the transmission coefficient  $T$ , is given by:

$$T = \frac{16}{|(1 + \frac{\alpha}{ik})(1 + \frac{ik}{\alpha}) \exp(-\alpha a) + (1 - \frac{\alpha}{ik})(1 - \frac{ik}{\alpha}) \exp(\alpha a)|^2},$$

where  $k = (2mE)^{1/2}/\hbar$  and  $\alpha = (2m(U_0 - E))^{1/2}/\hbar$ .

2. We consider a very large barrier where  $a \gg 1/\alpha$ , toward this limit, give the expression of  $T$  in function of  $E/U_0$  and  $\alpha a$ .
3. We consider the approximate formula:

$$\text{Log}(T) = -2\alpha a$$

What is the accuracy of this expression as compared to the result previously obtained in 2. If  $\alpha a = 50$ , show that the error is  $\simeq 1.4\%$ . What is happening at the classical limit (Hint: The planck constant goes to zero)?