

ECE609 Spring06
HOMEWORK 1
Review of Quantum Mechanics

1 The Photoelectric effect (15pts)

1. A metal has a workfunction of 4.3 eV. What is the minimum photon energy in Joule to emit an electron from this metal through the photo-electric effect? What are the photon frequency in Terahertz and the photon wavelength in micrometer (we note that $\lambda = c/\nu$ where c is the velocity of the light $\simeq 3 * 10^8 m.s^{-1}$)? What is the corresponding photon momentum ?
2. A photocathode in Potassium has a workfunction of 2 eV. What condition for the photon wavelength is necessary (in nanometer) to allow the photoelectric effect ?, For an incoming radiation of $\lambda = 546.1nm$, what is the maximum velocity of an extracted electron ? For an incoming radiation of $\lambda = 645.5nm$, what is the maximum velocity of an extracted electron ?

2 Hydrogen atom (30pts)

The fundamental level (1s state) of the hydrogen atom can be described by the wave function (spherical coordinates $r \geq 0$):

$$\Psi(r) = Cexp(-r/a_0),$$

where C is a constant (complex) and a_0 is defined as the Bohr radius.

(Hint: for all the integrations, we will consider a volume $4\pi r^2 dr$ - elementary volume between two spheres with radius r and $r + dr$ (shell))

1. Find the value of C ,
2. What is the density of probability to find an electron in a volume $4\pi r^2 dr$ (shell). Plot the density of probability dP/dr in function of r .
3. We call $r = r_p$ the value for which this probability exhibits a maximum. What is r_p in function of a_0 ?
Comment
4. What is the probability to find an electron at the distance between $0.9r_p$ and $1.1r_p$?
5. Give the average value (expectation value) of the distance that separate the proton from the electron $\langle r \rangle$. We note $\langle r \rangle = \langle \Psi | r | \Psi \rangle = \int_0^\infty \Psi^*(x)r\Psi(x)4\pi r^2 dr$.
6. Give the expectation values (in eV) of the potential energy $\langle U \rangle$ and the kinetic energy $\langle T \rangle$. What is the value of the total energy ? Comment.

3 Finite potential well (25pts)

We consider a 1D finite potential well, where the variations of $U(x)$ are defined by:

$$U(x) = \begin{cases} 0 & \text{if } x < -a/2 \text{ (region I), or } x > a/2 \text{ (region III)} \\ -U_0 & \text{if } -a/2 < x < a/2 \text{ (region II)} \end{cases}$$

where a and U_0 are positive constants. We consider an incoming electron wave from the left of the system with energy $E > 0$ and amplitude equal to one.

1. Give the expressions of the wave function $\Psi(x)$ in the three regions. We will set:

$$k = (2mE)^{1/2}/\hbar \text{ and } K = (2m(E + U_0))^{1/2}/\hbar.$$

2. Give the expression of the Transmission coefficient T . What is the reflection coefficient R (no calculation is necessary here)?
3. Show that T can take the following form:

$$T = \frac{1}{1 + f(E)\sin^2 g(E)}$$

where you will give the expressions of $f(E)$ and $g(E)$.

4. Numerical application, $a = 1.5A$, $U_0 = 16eV$. Plot $T(E)$ for $E < 150eV$
5. Calculate (analytically) the energy values for which T is equal to 1 (total transmission). What are the numerical expressions of these energy values (we consider only the values smaller than $150eV$) ?
6. What is happening if $E \rightarrow \infty$, Comment.

4 Finite potential wall (30pts)

We consider a 1D finite potential wall, where the variations of $U(x)$ are defined by:

$$U(x) = \begin{cases} 0 & \text{if } x \leq 0 \text{ (region I), or } x \geq a \text{ (region III)} \\ U_0 & \text{if } 0 < x < a \text{ (region II)} \end{cases}$$

where a and U_0 are positive constants.

We consider an incoming electron wave from the left of the system with energy $0 < E < U_0$ and amplitude equal to one.

1. Show that the transmission coefficient T , is given by:

$$T = \frac{16}{|(1 + \frac{\alpha}{ik})(1 + \frac{ik}{\alpha}) \exp(-\alpha a) + (1 - \frac{\alpha}{ik})(1 - \frac{ik}{\alpha}) \exp(\alpha a)|^2},$$

where $k = (2mE)^{1/2}/\hbar$ and $\alpha = (2m(U_0 - E))^{1/2}/\hbar$.

2. We consider a very large barrier where $a \gg 1/\alpha$, toward this limit, give the expression of T in function of E/U_0 and αa .
3. We consider the approximate formula:

$$\text{Log}(T) = -2\alpha a$$

What is the accuracy of this expression as compared to the result previously obtained in 2. If $\alpha a = 50$, show that the error is $\simeq 1.4\%$. What is happening at the classical limit (Hint: The planck constant goes to zero)?